

National Survey of Sea Lice (*Lepeophtheirus salmonis* Krøyer and *Caligus elongatus* Nordmann) on Fish Farms in Ireland - 2017

Pauline O'Donohoe

Frank Kane

Suzanne Kelly

Tom McDermott

Jack D'arcy

Joanne Casserly

Jamie Downes

Dave Jackson



Marine Institute
Foras na Mara

**NATIONAL SURVEY OF SEA LICE (*LEPEOPHTHEIRUS*
SALMONIS KRØYER AND *CALIGUS ELONGATUS* NORDMANN)
ON FISH FARMS IN IRELAND – 2017**

Irish Fisheries Bulletin No. 48

April 2018

Pauline O'Donohoe, Frank Kane, Suzanne Kelly, Tom McDermott,
Jack D'Arcy, Joanne Casserly, Jamie Downes and Dave Jackson.

Aquaculture Section,
Fisheries Ecosystem Advisory Services,
Marine Institute,
Rinville,
Oranmore,
Co. Galway.

www.marine.ie

<http://www.marine.ie/Home/site-area/areas-activity/aquaculture/sea-lice>

TABLE OF CONTENTS

Table of Contents	3
Figures.....	3
Tables.....	3
INTRODUCTION.....	1
History of Sea Lice Monitoring in Ireland	3
Sea Lice Control Methods.....	5
METHODOLOGY	6
RESULTS	8
Atlantic salmon 2016 (one-sea-winter salmon)	8
Atlantic salmon 2017 (smolts)	10
Sampling record	10
One-sea-winter salmon monthly trend by bay	10
Regional monthly means for one-sea-winter salmon	13
Annual trends	14
DISCUSSION	15
REFERENCES	17
APPENDIX I. Mean sea lice levels on salmonid farms in 2017.....	19

FIGURES

Figure 1. Life cycle of <i>Lepeophtheirus salmonis</i> (after Schram 1993 & Hamre 2013).	2
Figure 2. Locations of fish farm sites.	7
Figure 3. Mean (SE) ovigerous <i>L. salmonis</i> per fish per month per region in 2017 on one-sea-winter salmon fish... ..	13
Figure 4. Mean (SE) mobile <i>L. salmonis</i> per fish per month per region in 2017 on one-sea-winter salmon.....	13
Figure 5. Annual trend (May mean) (SE) ovigerous <i>L. salmonis</i> on one-sea-winter salmon.	14
Figure 6. Annual trend (May mean) (SE) mobile <i>L. salmonis</i> on one-sea-winter salmon.	14

TABLES

Table 1. Veterinary medicines authorized to assist in the control of sea lice on salmonids in Ireland	5
Table 2. Summary of inspections results on one-sea-winter salmon nationally in 2017.....	8
Table 3. Summary of inspections results on one-sea-winter salmon in the Southwest in 2017.....	8
Table 4. Summary of inspections results on one-sea-winter salmon in the West in 2017.....	9
Table 5. Summary of inspections results on one-sea-winter salmon in the Northwest in 2016.....	9
Table 6. Summary of inspections results on salmon smolts nationally in 2017.	10
Table 7. Mean ovigerous and mean mobile <i>Lepeophtheirus salmonis</i> and <i>Caligus elongatus</i> levels per month, for one-sea-winter salmon, for each bay inspected in the year 2017.....	12

INTRODUCTION

Sea lice are a naturally occurring parasite found on most fish, including salmonids. They are small ecto-parasitic copepod crustaceans which occur on several species of fish. There are estimated to be approximately 559 species made up of 37 genera (Ahyong et al., 2011), including 162 *Lepeophtheirus* species (Chad & Goeff, 2011) and 268 *Caligus* species (Boxshall, 2011). The two main species found in Ireland are *Caligus elongatus* and *Lepeophtheirus salmonis*. *L. salmonis* infests only salmonids, while *C. elongatus* is known to parasitise over 80 different species of marine fish. *L. salmonis* is the larger of the two species and is regarded as the more damaging parasite. It is endemic at a high prevalence (>90%) within wild populations (Jackson et al., 2013), and occurring frequently on farmed salmonids (Jackson & Minchin, 1992; Jackson et al., 2005). There are two species of salmonids farmed in Ireland on a commercial basis, Atlantic salmon *Salmo salar* L. and rainbow trout *Oncorhynchus mykiss* Walbaum.

L. salmonis is an obligate parasite with a direct lifecycle, with 8 stages, comprising of nauplius 1 and 2, copepodid, chalimus 1 and 2, preadult 1 and 2 and the adult stages. The nauplius 1 stage hatches from paired egg-strings and is dispersed in the plankton. It moults to nauplius 2, also planktonic, which is followed by a copepodid, the infective stage where attachment to the host takes place. The copepodid then moults through the attached chalimus stages before becoming a mobile pre-adult. There are two pre-adult stages before maturing to the adult phase. The adult female can produce a number of batches of paired egg-strings, which in turn hatch into the water column to give rise to the next generation (Hamre, 2013; Kabata, 1979; Schram, 1993). The mean length for an adult female is 8mm-11mm and an adult male is 5mm-6mm (Schram, 1993). Under experimental conditions female *L. salmonis* survived up to 210 days, producing as many as 11 pairs of egg strings (Boxaspen, 2006). Jackson and Minchin (1992), in Ireland, found fecundity (mean eggs per pair of egg strings) on wild salmon to be 965 ± 30 , which was higher than for farmed salmon at 758 ± 39 . This contrasts to a lower fecundity recorded for wild and farmed salmon in Norway where mean egg numbers have been recorded as 304 ± 32 with a range from 246 to 366 at 7.2°C (Heuch et al. 2000).




Figure 1. Life cycle of *Lepeophtheirus salmonis* (after Schram 1993 & Hamre 2013).

C. elongatus is smaller in size than *L. salmonis* averaging approximately 6-8mm in length (Hogans & Trudeau, 1989). The fact that *C. elongatus* is not as host specific as *L. salmonis* (Kabata, 1979) and that the hosts migrate widely is thought to be a factor in the highly variable levels on farmed salmonids at different times of the year.

History of Sea Lice Monitoring in Ireland

In 1991, the Department of the Marine instigated a Sea Lice Monitoring Programme for finfish farms in Ireland (Jackson & Minchin, 1993) and in 1993 monitoring was expanded nationwide (Jackson et al., 2002; Jackson et al., 2005). In May 2000 the protocol for sea lice monitoring was formally published (*Monitoring Protocol No.3 for Offshore Finfish Farms – Sea Lice Monitoring and Control*) by the Department of Marine and Natural Resources.

In 2008, the Department of Agriculture, Fisheries and Food (DAFF) published “A strategy for the improved pest control on Irish salmon farms”. The strategy outlines a comprehensive range of measures to provide for enhanced sea lice control and was developed by a joint DAFF, Marine Institute and Bord Iascaigh Mhara workgroup in response to difficulties experienced by farms in achieving the low levels of infestation required by the national control programme. These measures draw on the on-going Single Bay Management (SBM) process and, through a comprehensive action plan and list of recommendations, seek to advance the suite of tools necessary for improved sea lice control on farms.

The objectives of the National Sea Lice Monitoring Programme are:

- To provide an objective measurement of infestation levels on farms.
- To investigate the nature of infestations.
- To provide management information to drive the implementation of control and management strategies.
- To facilitate further development and refinement of this strategy.

The sea lice control and management strategy has five principal components:

- Separation of generations.
- Annual fallowing of sites.
- Early harvest of two-sea-winter fish.
- Targeted treatment regimes, including synchronous treatments.
- Agreed husbandry practices.

These components combine to reduce the development of sea lice infestations and to ensure the most effective treatment of sea lice. They seek to minimise sea lice levels whilst decreasing reliance on, and reducing the use of, veterinary medicines. Separation of generations and annual fallowing prevent the transmission of infestations from one generation to the next. The early harvest of two-sea-winter fish removes a potential reservoir of sea lice, while the agreed husbandry practises and targeted treatments enhance the efficacy of treatment regimes. One important aspect of targeted treatments is carrying out of synchronized autumn/winter treatments to reduce sea lice burdens to as close to zero as practicable, on all fish which are to be over-wintered. This is fundamental to

achieving near zero egg-bearing sea lice in spring. The agreed husbandry practises cover a range of related fish health, quality and environmental issues in addition to those specifically related to sea lice control. The Single Bay Management Programme serves to facilitate this and provides a forum for exchange of information between farmers.

In late winter and early spring seawater temperatures are at a minimum and the development rate of sea lice is slower. Rising water temperatures in spring tend to synchronise the development of sea lice larvae. A strategic treatment at this time can break the cycle of infection. Ovigerous female sea lice are those which produce the infective larvae and treatments are timed to remove adult females before they can release larvae. Setting the treatment trigger at 0.5 ovigerous *L. salmonis* per fish in spring ensures that treatments are carried out when a maximum of half of the fish examined have one ovigerous sea louse. This is a practical time to interrupt sea lice development. Later in the year, the development of new generations are not as synchronised and automatic intervention at a sea lice level of 0.5 ovigerous by way of treatment is not justified. A level of 2.0 ovigerous sea lice per fish has been shown to be a pragmatic level at which intervention by way of treatment is advisable. Levels of mobile and juvenile sea lice are important in advising fish health professionals in developing a sea lice control strategy. However, they are not of themselves appropriate measures upon which to trigger mandatory treatments.

Results of the monitoring programme are sent to the relevant farm within 5-10 days of each inspection. A monthly report of results is circulated to relevant parties and the data is published annually (www.marine.ie; Copley *et al.*, 2001; McCarney *et al.*, 2002; O'Donohoe *et al.*, 2003-2017).

Sea Lice Control Methods

The sea lice control and management strategy process includes the use of husbandry, management practises, and chemotherapeutants to control the numbers of sea lice on farms. Table I shows a list of the veterinary medicines authorised to assist in the control of sea lice in Ireland in 2016. These are either administered topically or incorporated into the diet. Topical treatments are administered by bathing the fish in specified concentrations of the medicine. Bath treatments can be conducted by the use of well-boats or tarpaulins/skirts to enclose the salmon net-pens. Medicines incorporated into the diet are a very efficient way to get the required dose to the fish. An over-reliance on any one technique can result in reduced efficacy in the short term and lead to development of resistance in time. A multi-pronged approach to sea lice control is considered more effective in the long-term.

The use of cleaner fish for the control of sea lice is being pursued in Ireland. Various wrasse species were used on salmon farms in the 1990s but recent interest has mainly been in stocking ballan wrasse *Labrus bergylta* (Treasurer, 2013). Studies in Norway also indicate that lumpfish *Cyclopterus lumpus* is a suitable cold-water option for biological delousing of Atlantic salmon (Imsland, 2014). Lumpfish are currently being trialled on farms in Ireland as part of a sea lice management plan, with positive results reported.

The use of filtration methods at harvest sites has also proven to be a very successful method for removing all stages of sea lice, including egg-strings; preventing sea lice from re-entering the water column and potentially re-infecting stocks adjacent to the harvest area (O'Donohoe & McDermott, 2014).

Table I. Veterinary medicines authorized to assist in the control of sea lice on salmonids in Ireland (www.hpra.ie).

Compound	Group	Licensing status	Delivery Method	Mode of action	Stages targeted	Withdrawal period
Animal medicines						
Deltamethrin	Pyrethroid	Full MA	Bath	Interferes with nerve transmission by blocking sodium channels in nerve cells	Adults, Preadults. Chalimus unknown	5 degree-days
Emamectin benzoate	Avermectin	Full MA	In-feed	Interferes with neurotransmission disrupting nerve cells causing paralysis and death	All stages	Zero
Hydrogen peroxide	Oxidizer	Full MA	Bath	Gas embolism	Adults, Preadults	Zero

MA - marketing authorisation from the Health Products Regulatory Authority.

METHODOLOGY

Farmed stocks of Atlantic salmon in Ireland are inspected on 14 occasions throughout the year to monitor sea lice levels as part of a national programme. Additional follow-up inspections may be carried out when it is deemed appropriate. Sea lice inspections take place twice per month in March, April and May (the spring period) and then monthly for the remainder of the year. December and January are combined and only one inspection is carried out.

At each inspection two samples are taken for each generation of fish on site, a sample from a standard cage, which is sampled at each subsequent inspection, and a sample from a random cage, which is chosen on the day of the inspection. Thirty fish are examined for each sample after anaesthetising using tricaine methane sulfonate in seawater. The seawater is sieved for any detached lice at the end of each sample. Each fish is examined individually for all mobile sea lice. Sea lice are removed and preserved in 70% ethanol. In the laboratory the species, quantity and life-stage of the sea lice are determined and recorded. The mean number of sea lice per fish is calculated (including the number of detached sea lice from the sieved seawater). Results presented are mean ovigerous sea lice levels and mean total mobile sea lice levels for *L. salmonis* and *C. elongatus* per fish.

Ovigerous sea lice levels are a measure of the breeding female population and total mobile levels measure successful infestation levels. The information gathered aims to evaluate the level of sea lice on the fish and to inform the farmer on a sea lice management strategy. Effective parasite control is characterised by a drop in sea lice levels on the subsequent inspection.

There are three distinct regions in Ireland where salmonid farming is carried out; the Southwest (Counties Cork and Kerry), the West (Counties Mayo and Galway) and the Northwest (Co. Donegal). These regions are geographically separate from each other with distances between regions of c.160 km from Northwest to West and c.200 km from West to Southwest (Figure 2).




Figure 2. Locations of fish farm sites.

RESULTS

During 2017 a total of 196 sea lice inspections were carried out on the 21 active farm sites. Over 90% of Atlantic salmon samples were below the Treatment Trigger Levels (TTL) as outlined in the *Monitoring Protocol No.3 for Offshore Finfish Farms – Sea Lice Monitoring and Control*, Department of Marine and Natural Resources (2000). Of the 101 inspections carried out on salmon smolts 97% were below the TTL and 85% of the 95 inspections carried out on one-sea-winter salmon were below TTL.

Results of monthly sea lice inspections of all active salmonid sites for 2017 are presented in Appendix I.

Atlantic salmon 2016 (one-sea-winter salmon)

One-sea-winter salmon were stocked in a total of 13 sites in 10 bays in 2017. Ninety-five visits were undertaken to this generation of fish.

Ovigerous *L. salmonis* levels greater than the TTL were recorded in a total of 15 inspections (16%) on one-sea-winter fish (Table 3). Within the critical spring period sea lice levels were in excess of 0.5 ovigerous females per fish on 11 inspections (19%) and outside of the spring period 4 inspections (11%) were in excess of 2.0 ovigerous female *L. salmonis* per fish.

Table 2. Summary of inspections results on one-sea-winter salmon nationally in 2017.

	Samples in Spring	Over in Spring	Samples outside Spring	Over outside Spring	Total Samples	Total Over	% over in Spring	% over outside Spring	% over total
National Totals	57	11	38	4	95	15	19%	11%	16%

C. elongatus levels greater than 10 individuals per fish were recorded on one occasion during the year.

Southwest Region

In the Southwest there were no recorded instances of *L. salmonis* levels greater than the treatment trigger levels (Table 4).

Table 3. Summary of inspections results on one-sea-winter salmon in the Southwest in 2017

Company	Site	Samples in Spring	Over in Spring	Samples outside Spring	Over outside Spring	Total Samples	Total Over	% over in Spring	% over outside Spring	% over total
Marine Harvest Ireland	Roanarraig- Ahabeg	1	0	1	0	2	0	0%	0%	0%
	Inishfarnard	6	0	4	0	10	0	0%	0%	0%
Southwest	Totals	7	0	5	0	12	0	0%	0%	0%

West Region

In the West, *L. salmonis* infestation levels greater than the TTL were recorded on 7 out of 24 inspections (29%) in the spring period and on 4 out of 18 inspections (22%) outside the spring period (Table 5).

Table 4. Summary of inspections results on one-sea-winter salmon in the West in 2017.

Company	Site	Samples in Spring	Over in Spring	Samples outside Spring	Over outside Spring	Total Samples	Total Over	% over in Spring	% over outside Spring	% over total
Bradán Beo Teo.	Ardmore	6	0	2	0	8	0	0%	0%	0%
Bifand Ltd.	Fraochoilean	5	1	4	2	9	3	0%	50%	33%
Comhlucht Bradáin Chonamara Teo.	Sealax	1	0	2	0	3	0	0%	0%	0%
Rosroe Salmon Ltd	Inishdeigil	6	2	2	0	8	2	33%	0%	25%
Clare Island Seafarms Ltd.	Portlea	6	4	8	2	14	6	67%	25%	43%
West	Totals	24	7	18	4	42	11	29%	22%	26%

Northwest Region

The treatment trigger levels were exceeded on 4 of the 26 inspections in the spring and none of the 15 inspections outside the spring period in the Northwest (Table 6).

Table 5. Summary of inspections results on one-sea-winter salmon in the Northwest in 2016.

Company	Site	Samples in Spring	Over in Spring	Samples outside Spring	Over outside Spring	Total Samples	Total Over	% over in Spring	% over outside Spring	% over total
Ocean Farm Ltd.	Ocean Inver	6	1	2	0	8	1	17%	0%	13%
Marine Harvest Ireland	Creevin	3	3	2	0	5	3	100%	0%	60%
	Cranford B	5	0	1	0	6	0	0%	0%	0%
	Glinsk	0	0	1	0	1	0	0%	0%	0%
	Millstone	6	0	3	0	9	0	0%	0%	0%
	Lough Swilly	6	0	6	0	12	0	0%	0%	0%
Northwest	Totals	26	4	15	0	41	4	15%	0%	10%

Atlantic salmon 2017 (smolts)

A total of 101 inspections were made to 11 sites stocking Atlantic salmon 2017 S1 and S½ smolts during the year 2017. *L. salmonis* levels were below the TTL of ovigerous female lice per fish for all of the inspections (100%) carried out in spring and were over for 3 inspections outside the spring period (Table 7).

Table 6. Summary of inspections results on salmon smolts nationally in 2017.

Company	Site	Samples in Spring	Over in Spring	Samples outside Spring	Over outside Spring	Total Samples	Total Over	% over in Spring	% over outside Spring	% over total
Marine Harvest Ireland	Roanarraig-Ahabeg	6	0	8	0	14	0	0%	0%	0%
	Deenish	2	0	6	0	8	0	0%	0%	0%
Southwest	Totals	8	0	14	0	22	0	0%	0%	0%
Bradán Beo Teo.	Daonish	6	0	8	0	14	0	0%	0%	0%
Comhlucht Bradáin Chonamara Teo.	Outer Bertraghboy Bay	1	0	6	0	7	0	0%	0%	0%
Mannin Bay Salmon Company Ltd	Hawk's Nest	6	0	6	0	12	0	0%	0%	0%
Rosroe Salmon Ltd.	Inishdeighil	0	0	2	1	2	1	0%	50%	50%
Clare Island Seafarms Ltd	Clare Island Portlea	2	0	4	0	6	0	0%	0%	0%
	Seastream	0	0	2	0	2	0	100%	0%	0%
Curraun Blue Ltd	Curraun	6	0	8	1	14	1	0%	13%	7%
West	Totals	21	0	36	2	57	2	0%	6%	4%
Ocean Farm Ltd.	Mc Swynes	6	0	8	1	14	1	0%	13%	7%
Marine Harvest Ireland	Glinsk	2	0	6	0	8	0	0%	0%	0%
Northwest	Totals	8	0	14	1	22	1	0%	7%	5%
National Totals		37	0	64	3	101	3	0%	5%	3%

The maximum *C. elongatus* level recorded was 16.34 in the West in November in 2017.

Sampling record

Three samples were missed due to fish health issues in the 2017 sampling year (Roanarraig-Ahabeg in March and in both inspections in April).

One-sea-winter salmon monthly trend by bay

Bay mean ovigerous and mean mobile *L. salmonis* and *C. elongatus* levels for each bay are shown in Table 7 for one-sea-winter salmon throughout the year. Monthly ovigerous *L. salmonis* levels greater than the spring TTL of 0.5 ovigerous sea lice per fish were recorded on 7 of the 26 occasions. Of these, 2 were in Clew Bay, Killary Harbour and Donegal Bay, and one in Ballinakill Bay.

Bay mean ovigerous levels of 2.0 ovigerous females per fish or greater were recorded on 4 out of 34 occasions, outside of the spring period. These occurred on 2 occasions at both Clew Bay and Ballinakill Harbour.

Mean mobile levels per bay in excess of 10 *L. salmonis* per fish were recorded on 6 occasions, 2 of these instances had means of greater than 20 mobile lice per fish. The maximum bay mean level recorded was 53.28 mobile sea lice per fish, in Ballinakill Harbour in September.

Table 7. Mean ovigerous and mean mobile *Lepeophtheirus salmonis* and *Caligus elongatus* levels per month, for one-sea-winter salmon, for each bay inspected in the year 2017.

Mean ovigerous <i>L. salmonis</i>											
	Dec/ Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov
Bantry Bay	0.12		0.20								
Kenmare Bay	0.00	0.03	0.12	0.23	0.19	0.52	0.07	HO			
Kilkieran Bay	0.02	0.10	0.15	0.03	0.01	HO					
Bertraghboy Bay	0.84	0.30	0.21	TO							
Ballinakill Harbour			0.08	0.23	0.58	0.61	1.22	3.78	15.21	HO	
Killary Harbour	0.63	0.79	0.78	0.35	0.57	HO					
Clew Bay	0.00	0.70	0.35	1.44	1.96	1.18	0.60	1.00	1.46	3.85	8.70
Donegal Bay	0.66	1.13	0.70	0.52	0.09	HO					
Mulroy Bay	0.09	0.33	0.18	0.02	0.11	0.10	HO				
Lough Swilly		0.04	0.04	0.00	0.09	0.27	1.47	1.02	0.00	0.02	HO
Mean mobile <i>L. salmonis</i>											
	Dec/ Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov
Bantry Bay	1.20		0.40								
Kenmare Bay	0.09	0.17	0.45	1.63	4.11	11.54	0.36	HO			
Kilkieran Bay	0.44	2.49	1.31	0.51	0.42	HO					
Bertraghboy Bay	4.82	1.02	0.61	TO							
Ballinakill Harbour			0.23	0.58	2.94	1.13	8.07	16.26	53.28	HO	
Killary Harbour	2.71	3.09	4.02	6.06	5.49	HO					
Clew Bay	1.13	3.20	3.41	7.69	10.28	4.36	3.54	5.19	9.53	9.47	45.04
Donegal Bay	7.64	3.68	4.28	9.86	1.06	HO					
Mulroy Bay	0.29	0.99	0.46	0.29	0.25	0.90	HO				
Lough Swilly		0.15	0.16	0.11	0.33	1.39	4.77	15.60	1.19	0.38	HO
Mean ovigerous <i>C. elongatus</i>											
	Dec/ Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov
Bantry Bay	0.57		0.20								
Kenmare Bay	0.72	0.03	0.35	0.13	0.20	0.07	0.00	HO			
Kilkieran Bay	1.99	1.18	0.03	0.00	0.07	HO					
Bertraghboy Bay	0.04	0.00	0.00	TO							
Ballinakill Harbour			0.00	0.01	0.00	0.00	0.02	0.00	0.04	HO	
Killary Harbour	0.22	0.20	0.00	0.00	0.00	HO					
Clew Bay	1.83	1.80	3.16	0.64	0.40	0.04	0.06	0.07	0.23	0.09	1.45
Donegal Bay	0.90	0.02	0.02	0.07	0.00	HO					
Mulroy Bay	0.34	0.08	0.00	0.00	0.00	0.00	HO				
Lough Swilly		0.04	0.00	0.04	0.00	0.08	0.00	0.12	0.07	0.29	HO
Mean mobile <i>C. elongatus</i>											
	Dec/ Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov
Bantry Bay	1.48		0.30								
Kenmare Bay	1.10	0.03	0.69	0.29	0.41	0.27	0.04	HO			
Kilkieran Bay	3.98	2.33	0.05	0.04	0.13	HO					
Bertraghboy Bay	0.41	0.07	0.00	TO							
Ballinakill Harbour			0.00	0.02	0.01	0.00	0.02	0.00	0.12	HO	
Killary Harbour	0.54	0.60	0.10	0.00	0.01	HO					
Clew Bay	5.49	3.90	8.35	1.26	0.57	0.07	0.27	0.21	0.53	0.16	2.58
Donegal Bay	3.34	0.22	0.14	0.16	0.00	HO					
Mulroy Bay	1.10	0.19	0.00	0.00	0.01	0.03	HO				
Lough Swilly		0.18	0.05	0.07	0.00	0.15	0.00	0.14	0.56	0.72	HO

HO = Harvested out

Regional monthly means for one-sea-winter salmon

L. salmonis ovigerous and monthly mean mobiles levels per fish for one-sea-winter salmon regionally are shown in Figures 3 and 4. In the spring period of 2017 the ovigerous mean sea lice levels per fish exceeded TTLs for March, April and May in the West, and for April in the Northwest.




Figure 3. Mean (SE) ovigerous *L. salmonis* per fish per month per region in 2017 on one-sea-winter salmon fish.




Figure 4. Mean (SE) mobile *L. salmonis* per fish per month per region in 2017 on one-sea-winter salmon.

Total mobile *L. salmonis* levels exceeded 10 sea lice per fish in August, September and November in the West, and once in both of the Southwest and Northwest regions. Total regional mean mobile *L. salmonis* levels peaked at 11.54 mobile sea lice per fish in the Southwest, 45.04 mobile sea lice per fish in the West and at 15.60 mobile sea lice per fish in the Northwest.

Annual trends

The annual trends of *L. salmonis* ovigerous and mobile sea lice levels are compared in Figures 5 and 6 for one-sea-winter salmon in the month of May from 1991 to 2017.




Figure 5. Annual trend (May mean) (SE) ovigerous *L. salmonis* on one-sea-winter salmon.




Figure 6. Annual trend (May mean) (SE) mobile *L. salmonis* on one-sea-winter salmon.

Mean ovigerous *L. salmonis* levels in May of 2017 decreased to 0.43 sea lice per fish, compared to 0.45 in 2016. Total mobile levels of 2.86 showed a slight increase compared to May 2016 at 2.52 sea lice per fish.

DISCUSSION

Sea lice levels on smolts in 2017 were low. All of the sea lice inspections carried out on smolts were below the Treatment Trigger Levels (TTL); These results are similar to 2016.

In 2017 84% of inspections of one-sea-winter salmon were below TTL. During the spring period, in the Northwest, 85% of inspections were below the TTL compared to 91% in 2016. In the West, for the same period, 71% were below TTL which is a decrease from 63% on 2016. The Southwest again continued to have no breaches of protocol levels in 2017. All inspections of one-sea-winter salmon undertaken outside of the spring period, were below TTL in the Northwest and Southwest, while 78% were below in the West.

Sea lice levels in excess of 10 *L. salmonis* mobiles per fish on one-sea-winter salmon nationally were recorded on 13 occasions, compared to 8 occasions in 2016 and 17 in 2015, two of these inspections had means of greater than 20 mobile *L. salmonis* per fish. The highest mean sea lice level recorded for one-sea-winter salmon was 53.28 mobile *L. salmonis* per fish, this compares to 37.6 mobile *L. salmonis* per fish in 2016 and 71.5 mobile *L. salmonis* per fish in 2015. There were no unusually high numbers of *Caligus elongatus* recorded in 2017.

Sea lice levels across the country remained similar to previous years. Infestation on smolts was low as is the common annual trend. Sea lice levels on grower fish continued to show a level of control in line with annual trends over the last number of years. Higher sea lice levels were observed in the run up to harvest on larger grower fish which is in keeping with recorded data from sea lice inspections of harvest fish. Control methods including sea lice chemotherapeutants, mechanical and biological delousing methods and Single Bay Management practices continue to have a positive impact on the control of sea lice in Irish salmon farming.

GLOSSARY

<i>Grower:</i>	A fish which has been at sea for one complete year or longer.
<i>Mobile lice:</i>	All sea lice that are mobile – male and female (pre-adult and adult stages) sea lice that have developed beyond the attached larval stages.
<i>n<10</i>	Ten fish or fewer were inspected in one or both cages sampled.
<i>Ovigerous lice:</i>	An egg bearing adult female sea lice.
<i>Random (Ran.) Cage:</i>	A cage which is selected by the Inspector on the day of inspection.
<i>Salmonids:</i>	A fish of the family Salmonidae. It includes salmon, trout and charr.
<i>Standard (Std.) Cage:</i>	The selected cage which is sampled at each inspection.
<i>S1 Smolt:</i>	This pertains to a stage in the life cycle of the salmon when it changes from being a freshwater fish to a seawater fish, a process known as smoltification. These fish are transported to the saltwater environment in the spring, which is approximately 15 months after they were hatched.
<i>S½ Smolt (also known as S0):</i>	These fish are put under lights to hasten the onset of smoltification. An S½ smolt is ready to go to sea during the autumn/winter, approximately 11 months after hatching. They are sometimes referred to as S0 (S zero) smolts.
<i>SE:</i>	Standard error (error bars in the graphs) is the standard error of the mean of a sample from a population with a normal distribution, which is equal to the standard deviation of the normal distribution divided by the square root of the sample size.

REFERENCES

- Ahyong ST, Lowry JK, Alonso M, Bamber RN, Boxshall GA, Castro P, Gerken S, Karaman GS, Goy JW, Jones DS, Meland K, Rogers DC & Svavarsson J 2011. In: *Animal biodiversity: an outline of higher-level classification and survey of taxonomic richness*. Ed; Z-Q Zhang. *Zootaxa* 3148: 165–191.
- Boxaspen, K 2006. A review of the biology and genetics of sea lice. *ICES Journal of Marine Science*, 63, 1304–1316.
- Boxshall G 2011. "*Caligus* O.F. Müller, 1785". In: World Copepoda database. *World Register of Marine Species*. TC Walter & G Boxshall.
- Chad WT & Geoff G 2011. "*Lepeophtheirus* von Nordmann, 1832". In T. C. Walter & G. Boxshall. *World Copepoda database*. *World Register of Marine Species*.
- Copley L, McCarney P, Jackson D, Hasset D, Kennedy S & Nulty C 2001. The occurrence of Sea lice (*Lepeophtheirus salmonis* Krøyer) on Farmed Salmon in Ireland (1995 to 2000). *Marine Resource Series*, No.17. Marine Institute.
- Department of Agriculture, Fisheries and Food 2008. A strategy for the improved pest control on Irish salmon farms.
- Department of the Marine and Natural Resources 2000. Monitoring Protocol No. 3: Offshore Finfish Farms - Sea Lice Monitoring and Control.
- Hamre LA, Eichner C, Caipang CMA, Dalvin ST, Bron JE, Nilsen F, Boxshall G, Skern-Mauritzen R 2013. The Salmon Louse *Lepeophtheirus salmonis* (Copepoda: Caligidae) Life Cycle Has Only Two Chalimus Stages. *PLoS ONE* 8(9): e73539. doi:10.1371/journal.pone.0073539
- Hogans WE & DJ Trudeau 1989. Preliminary studies on the biology of sea lice, *Caligus elongatus*, *Caligus curtus* and *Lepeophtheirus salmonis* (Copepoda: Caligida) parasitic on cage-cultured salmonids in the Lower Bay of Fundy. *Canadian Technical Report of Fisheries and Aquatic Sciences*, No. 1715: 14 pp.
- Heuch PA, Nordhagen JR, Schram TA 2000. Egg production in the salmon louse [*Lepeophtheirus salmonis* (Krøyer)] in relation to origin and water temperature. *Aquaculture Research* 31:805-814.
- Imsland AK, Reynolds P, Eliassen G, Hangstad TA, Foss A, Vikingstad E, Elvegård TA 2014. The use of lumpfish (*Cyclopterus lumpus* L.) to control sea lice (*Lepeophtheirus salmonis* Krøyer) infestations in intensively farmed Atlantic salmon (*Salmo salar* L.) *Aquaculture* 424–425 (2014) 18–23.
- Jackson D, Copley L, Kane F, Naughton O, Kennedy S & O'Donohoe P 2005. Thirteen years of monitoring sea lice in farmed salmonids. In 'Long-term monitoring: Why, What, Where, When & How?' Edited by John Solbé, *Proceedings of 'The importance of long-term monitoring of the environment' conference*. Sherkin Island Marine Station, Ireland. pp 92-105.
- Jackson D & Minchin D 1992. Aspects of the reproductive output of two caligid copepod species on cultivated salmon. *Invertebrate Reproductive Development*, 22: 87-90.
- Jackson D & Minchin D 1993. Lice infestations of farmed salmon in Ireland. In: *Pathogens of wild and farmed fish* Ed; GA Boxshall & D Defaye, Ellis Horwood Ltd, London, UK. pp. 188–201.
- Jackson D, Hasset D & Copley L 2002. Integrated lice management on Irish Salmon farms. *Fish Veterinary Journal*, (6) 28-38.
- Jackson D, Kane F, O'Donohoe P, Mc Dermott T, Kelly S, Drumm A & Newell J 2013. Sea lice levels on wild Atlantic salmon, *Salmo salar* L., returning to the coast of Ireland. *Journal of Fish Diseases*, 36: 293–298. doi: 10.1111/jfd.12059
- Kabata Z 1979. *Parasitic Copepoda of British Fishes*. The Ray Society, London.
- McCarney P, Copley L, Kennedy S, Nulty C & Jackson D 2002. National Survey of Sea lice (*Lepeophtheirus salmonis* Krøyer and *Caligus elongatus* Nordmann) on Fish Farms in Ireland – 2000 & 2001. *Fisheries Leaflets 180 & 181*, Marine Institute.

- O'Donohoe, et al.** 2017. National Survey of Sea lice (*Lepeophtheirus salmonis* Krøyer and *Caligus elongatus* Nordmann) on Fish Farms in Ireland. Published annually in *Irish Fisheries Bulletins Numbers 22, 24, 28, 31, 32, 33, 34, 40, 41, 44, 45, 46, 47 and Fisheries Leaflets 183 & 184*. Marine Institute.
- O'Donohoe P & McDermott T** 2014. Reducing sea lice re-infestation risk from harvest water at a salmon farm site in Ireland using a bespoke sieving and filtration system. *Aquacultural Engineering*. Volume 60, May 2014, Pages 73–76. <http://dx.doi.org/10.1016/j.aquaeng.2014.04.004>
- Schram TA** 1993. Supplementary descriptions of the developmental stages of *Lepeophtheirus salmonis* (Krøyer, 1837) (Copepoda: Caligidae). In: *Pathogens of wild and farmed fish* Ed; GA Boxshall & D Defaye, Ellis Horwood Ltd, London, UK. pp. 30-47.
- Treasurer J** 2013. Use of wrasse in sea lice control. *Scottish Aquaculture Resource Forum*. ISBN 978-1-907266-56-0

APPENDIX 1. MEAN SEA LICE LEVELS ON SALMONID FARMS IN 2017.

	Date	<i>Lepeophtheirus salmonis</i>		<i>Caligus elongatus</i>	
		F + eggs	Total	F + eggs	Total
BANTRY BAY					
MARINE HARVEST IRL.					
Roancharraig-Ahabeg					
Atlantic Salmon, 2016 S1/2	17/01/2017	0.12	1.20	0.57	1.48
	16/02/2017	On starve for harvest			
	09/03/2017	0.20	0.40	0.20	0.30
	29/03/2017	Not sampled for fish health reasons			
	April 2017	Not sampled for fish health reasons			
		Harvested Out			
Atlantic Salmon, 2017 S1/2	17/01/2017	0.00	0.04	0.02	0.05
	16/02/2017	0.02	0.25	0.70	1.37
	09/03/2017	0.11	0.23	0.00	0.00
	29/03/2017	0.11	0.38	0.00	0.00
	11/04/2017	0.09	0.24	0.00	0.00
	18/04/2017	0.08	0.56	0.02	0.08
	04/05/2017	0.09	1.26	0.02	0.02
	30/05/2017	0.10	0.76	0.00	0.02
	13/06/2017	0.19	0.68	0.00	0.03
	25/07/2017	0.05	0.33	0.29	0.42
	14/08/2017	0.07	0.23	0.75	1.49
	25/09/2017	0.03	0.08	0.00	0.00
	27/10/2017	0.02	0.04	1.72	3.44
	02/11/2017	0.00	0.02	1.82	3.59

	Date	<i>Lepeophtheirus salmonis</i>		<i>Caligus elongatus</i>	
		F + eggs	Total	F + eggs	Total
KENMARE BAY					
Deenish					
Atlantic Salmon, 2015 S1/2				Harvested Out	
Atlantic Salmon, 2017	04/05/2017	0.00	0.29	0.01	0.10
	30/05/2017	0.00	0.28	0.02	0.02
	13/06/2017	0.00	0.13	0.00	0.01
	26/07/2017	0.00	0.02	0.15	0.27
	15/08/2017	0.00	0.18	0.38	0.84
	25/09/2017	0.00	0.05	0.08	0.27
	27/10/2017	0.07	0.09	2.63	5.37
	02/11/2017	0.09	0.27	4.41	7.73
Inishfarnard					
Atlantic Salmon, 2016	17/01/2017	0.00	0.09	0.72	1.10
	16/02/2017	0.03	0.17	0.03	0.03
	09/03/2017	0.07	0.30	0.12	0.37
	29/03/2017	0.18	0.61	0.58	1.00
	11/04/2017	0.19	1.23	0.10	0.25
	18/04/2017	0.27	2.03	0.16	0.34
	04/05/2017	0.28	7.62	0.40	0.82
	30/05/2017	0.11	0.61	0.00	0.00
	13/06/2017	0.52	11.54	0.07	0.27
	25/07/2017	0.07	0.36	0.00	0.04
				Harvested Out	

	Date	<i>Lepeophtheirus salmonis</i>		<i>Caligus elongatus</i>	
		F + eggs	Total	F + eggs	Total
KILKIERAN BAY					
BRADAN BEO TEO.					
Ardmore					
Atlantic Salmon, 2016 S1/2	10/01/2017	0.02	0.44	1.99	3.98
	14/02/2017	0.10	2.49	1.18	2.33
	14/03/2017	0.29	2.42	0.02	0.05
	28/03/2017	0.00	0.20	0.05	0.05
	07/04/2017	0.00	0.23	0.00	0.06
	21/04/2017	0.06	0.79	0.00	0.02
	03/05/2017	0.00	0.23	0.04	0.07
	19/05/2017	0.02	0.61	0.10	0.19
Harvested Out					
Daonish					
Atlantic Salmon, 2017 S1/2	16/01/2017	0.03	0.13	0.00	0.03
	14/02/2017	0.00	1.82	0.00	0.02
	10/03/2017	0.17	1.76	0.04	0.20
	24/03/2017	0.00	0.21	0.00	0.00
	07/04/2017	0.08	0.38	0.00	0.02
	25/04/2017	0.02	0.24	0.00	0.00
	03/05/2017	0.03	0.88	0.02	0.07
	25/05/2017	0.12	0.83	0.07	0.15
	09/06/2017	0.02	0.16	0.00	0.02
	06/07/2017	0.33	1.33	0.40	0.57
	04/08/2017	0.05	0.36	0.02	0.12
	13/09/2017	0.12	0.67	0.05	0.07
	20/10/2017	0.00	0.14	0.07	0.08
	15/11/2017	0.00	0.07	0.08	0.20

	Date	<i>Lepeophtheirus salmonis</i>		<i>Caligus elongatus</i>	
		F + eggs	Total	F + eggs	Total
BERTRAGHBOY BAY					
BIFAND LTD. / MARINE HARVEST IRL.					
Outer Bertraghboy Bay					
Atlantic Salmon, 2017	19/05/2017	0.00	0.02	0.00	0.03
	07/06/2017	0.00	0.00	0.00	0.07
	04/07/2017	0.00	0.04	0.16	0.24
	25/08/2017	0.00	0.01	0.04	0.09
	21/09/2017	0.00	0.06	0.02	0.02
	26/10/2017	0.00	0.04	0.22	0.45
	13/11/2017	0.04	0.04	1.01	1.47
COMHLUCHT BRADAIN CHONAMARA TEO					
Sealax					
Atlantic Salmon, 2016	08/12/2016	0.84	4.82	0.04	0.41
	03/02/2017	0.30	1.02	0.00	0.07
	06/03/2017	0.21	0.61	0.00	0.00
Transferred to Fraochoilean					
CLIFDEN BAY					
MANNIN BAY SALMON COMPANY LTD.					
Hawks Nest					
Atlantic Salmon, 2017 S1/2	18/01/2017	0.00	0.00	0.27	0.36
	15/02/2017	0.00	0.06	0.48	0.73
	03/03/2017	0.00	0.24	0.70	1.08
	21/03/2017	0.05	0.33	0.41	0.43
	05/04/2017	0.10	0.34	0.03	0.09
	19/04/2017	0.07	0.21	0.00	0.00
	12/05/2017	0.05	0.29	0.00	0.00
	17/05/2017	0.00	0.42	0.00	0.00
	01/06/2017	0.10	0.45	0.00	0.00
	05/07/2017	0.12	4.45	0.02	0.05
Transferred to Inishdeighil					
Atlantic Salmon, 2017 S1/2	27/10/2017	1.03	4.87	0.00	0.00
	30/11/2017	0.34	1.34	0.00	0.00

		Date	<i>Lepeophtheirus salmonis</i>		<i>Caligus elongatus</i>	
			F + eggs	Total	F + eggs	Total
BALLINAKILL HARBOUR		<i>BIFAND LTD.</i>				
Fraochoilean						
Atlantic Salmon, 2015		Harvested Out				
Atlantic Salmon, 2016		31/03/2017	0.08	0.23	0.00	0.00
		06/04/2017	0.03	0.26	0.00	0.00
		18/04/2017	0.44	0.90	0.02	0.03
		08/05/2017	0.49	3.92	0.00	0.02
		26/05/2017	0.67	1.95	0.00	0.00
		01/06/2017	0.61	1.13	0.00	0.00
		13/07/2017	1.22	8.07	0.02	0.02
		23/08/2017	3.78	16.26	0.00	0.00
		26/09/2017	15.21	53.28	0.04	0.12
		Harvested out				
KILLARY HARBOUR						
<i>ROSROE SALMON LTD.</i>						
Inishdeighil						
Atlantic Salmon, 2016		16/02/2017	0.58	3.77	0.27	0.96
		13/03/2017	0.61	4.87	0.00	0.19
		31/03/2017	Sampled Atlantic Salmon, 2016 S1/2			
		13/04/2017	0.40	1.20	0.00	0.00
		28/04/2017	0.53	8.33	0.00	0.00
		09/05/2017	0.42	6.60	0.00	0.00
		23/05/2017	0.87	3.27	0.00	0.03
		Harvested Out				
Atlantic Salmon, 2016 S1/2		20/01/2017	0.63	2.71	0.22	0.54
		16/02/2017	1.00	2.41	0.12	0.24
		13/03/2017	1.48	6.83	0.00	0.16
		31/03/2017	0.17	0.79	0.00	0.00
		13/04/2017	0.04	2.75	0.00	0.00
		28/04/2017	0.42	11.97	0.00	0.00
		Harvested Out				
Atlantic Salmon, 2017 S1/2		22/08/2017	1.90	5.27	0.20	0.49
		25/09/2017	5.35	17.35	0.50	0.90
		Harvested and transferred to Hawks Nest				

	Date	<i>Lepeophtheirus salmonis</i>		<i>Caligus elongatus</i>	
		F + eggs	Total	F + eggs	Total
CLEW BAY					
CLARE ISLAND SEAFARMS LTD.					
Clare Island Smolt Site					
Atlantic Salmon, 2015				Harvested Out	
Atlantic Salmon, 2017	22/05/2017	0.00	0.19	0.05	0.07
	31/05/2017	0.00	0.16	0.00	0.00
	27/06/2017	0.06	0.43	0.37	0.61
	18/07/2017	0.00	0.11	0.02	0.09
	14/08/2017	0.00	0.24	0.02	0.43
	18/09/2017	0.00	0.12	0.00	0.00
			Transferred to Seastream		
Portlea					
Atlantic Salmon, 2016	06/01/2017	0.00	1.13	1.83	5.49
	22/02/2017	0.70	3.20	1.80	3.90
	08/03/2017	0.35	4.36	6.09	16.34
	27/03/2017	0.36	2.45	0.23	0.36
	11/04/2017	1.54	5.48	0.65	1.17
	26/04/2017	1.35	9.89	0.64	1.35
	22/05/2017	2.21	14.92	0.58	0.74
	31/05/2017	1.70	5.64	0.22	0.40
	27/06/2017	1.18	4.36	0.04	0.07
	18/07/2017	0.60	3.54	0.06	0.27
	14/08/2017	1.00	5.19	0.07	0.21
	18/09/2017	1.46	9.53	0.23	0.53
	19/10/2017	3.85	9.47	0.09	0.16
	22/11/2017	8.70	45.04	1.45	2.58
Seastream					
Atlantic Salmon, 2017	19/10/2017	0.00	0.08	0.07	0.07
	22/11/2017	0.08	11.21	3.53	9.58

		Date	<i>Lepeophtheirus salmonis</i>		<i>Caligus elongatus</i>		
			F + eggs	Total	F + eggs	Total	
BEALACRAGHER BAY	<i>CURRAUN BLUE LTD.</i>						
Curraun							
Atlantic Salmon, 2017 S1/2	06/01/2017	0.00	0.00	0.20	0.23	n<10	
	22/02/2017	0.00	0.12	0.46	0.71		
	08/03/2017	0.00	0.05	0.36	0.49		
	27/03/2017	0.00	0.02	0.32	0.37		
	11/04/2017	0.02	0.08	0.45	0.54		
	26/04/2017	0.00	0.12	0.68	1.29		
	18/05/2017	0.00	0.52	1.64	4.82		
	31/05/2017	0.14	0.97	0.00	0.05		
	27/06/2017	0.05	0.20	0.00	0.00		
	18/07/2017	0.96	3.30	0.00	0.00		
	16/08/2017	0.85	1.81	0.00	0.00		
	28/09/2017	0.65	1.67	0.00	0.00		
	25/10/2017	2.20	4.97	0.02	0.04		
	29/11/2017	1.21	6.26	0.00	0.00		

	Date	<i>Lepeophtheirus salmonis</i>		<i>Caligus elongatus</i>	
		F + eggs	Total	F + eggs	Total
DONEGAL BAY					
MARINE HARVEST IRL.					
Creevin					
Atlantic Salmon, 2016 S1/2	06/01/2017	0.14	8.18	0.93	3.05
	28/02/2017	1.87	4.93	0.00	0.37
	15/03/2017	0.80	3.62	0.00	0.02
	29/03/2017	0.78	6.22	0.00	0.04
	18/04/2017	1.08	15.30	0.00	0.04
Harvested Out					
OCEAN FARM LTD.					
Mc Swynes					
Atlantic Salmon, 2017 S1/2	06/01/2017	0.00	0.06	0.18	0.94
	15/02/2017	0.09	1.56	2.53	6.35
	06/03/2017	0.12	4.52	2.05	6.38
	28/03/2017	0.16	2.53	0.72	2.31
	05/04/2017	0.04	4.63	0.83	2.48
	19/04/2017	0.40	10.32	0.54	1.27
	09/05/2017	0.04	1.61	0.68	1.19
	29/05/2017	0.00	0.20	0.00	0.00
	15/06/2017	0.12	0.85	0.09	0.12
	13/07/2017	0.18	0.49	0.00	0.00
	16/08/2017	0.72	1.32	0.00	0.02
	19/09/2017	1.50	5.76	0.02	0.02
	25/10/2017	0.90	5.53	0.00	0.02
	15/11/2017	2.17	13.81	0.08	0.23

	Date	<i>Lepeophtheirus salmonis</i>		<i>Caligus elongatus</i>	
		F + eggs	Total	F + eggs	Total
Ocean Inver					
Atlantic Salmon, 2016 S1/2	06/01/2017	1.18	7.09	0.88	3.63
	15/02/2017	0.77	3.05	0.03	0.15
	06/03/2017	1.05	4.52	0.07	0.45
	28/03/2017	0.15	2.75	0.02	0.04
	04/04/2017	0.17	3.38	0.12	0.18
	18/04/2017	0.32	10.91	0.10	0.26
	09/05/2017	0.07	0.89	0.00	0.00
	29/05/2017	0.11	1.24	0.00	0.00
Harvested Out					
MULROY BAY					
MARINE HARVEST IRL.					
Cranford B					
Atlantic Salmon, 2016	16/02/2017	0.42	1.24	0.07	0.25
	07/03/2017	0.40	1.18	0.00	0.00
	21/03/2017	0.02	0.07	0.00	0.00
	05/04/2017	0.04	0.07	0.00	0.00
	19/04/2017	0.02	0.79	0.00	0.00
	10/05/2017	0.13	0.20	0.00	0.00
Harvested Out					
Glinsk					
Atlantic Salmon, 2016	05/01/2017	0.00	0.12	0.53	1.42
	Transferred to Lough Swilly				
Atlantic Salmon, 2017	10/05/2017	0.00	0.04	0.02	0.04
	30/05/2017	0.00	0.00	0.04	0.07
	16/06/2017	0.00	0.14	0.23	0.29
	12/07/2017	0.02	0.07	0.02	0.02
	17/08/2017	0.00	0.00	0.00	0.02
	20/09/2017	0.00	0.02	0.02	0.02
	06/10/2017	0.00	0.00	0.00	0.02
	15/11/2017	0.00	0.00	0.00	0.00

	Date	<i>Lepeophtheirus salmonis</i>		<i>Caligus elongatus</i>	
		F + eggs	Total	F + eggs	Total
Millstone					
Atlantic Salmon, 2016	05/01/2017	0.18	0.46	0.15	0.78
	16/02/2017	0.24	0.74	0.09	0.14
	07/03/2017	0.27	0.56	0.00	0.00
	21/03/2017	0.02	0.06	0.00	0.00
	06/04/2017	0.03	0.03	0.00	0.00
	20/04/2017	0.00	0.29	0.00	0.00
	10/05/2017	0.04	0.15	0.00	0.02
	30/05/2017	0.17	0.39	0.00	0.02
	16/06/2017	0.10	0.90	0.00	0.03
Harvested Out					
LOUGH SWILLY					
Lough Swilly					
Atlantic Salmon, 2015	Harvested Out				
Atlantic Salmon, 2016	16/02/2017	0.04	0.15	0.04	0.18
	07/03/2017	0.03	0.15	0.00	0.10
	21/03/2017	0.05	0.17	0.00	0.00
	06/04/2017	0.00	0.11	0.07	0.14
	20/04/2017	0.00	0.12	0.00	0.00
	10/05/2017	0.04	0.22	0.00	0.00
	30/05/2017	0.15	0.45	0.00	0.00
	16/06/2017	0.27	1.39	0.08	0.15
	12/07/2017	1.47	4.77	0.00	0.00
	17/08/2017	1.02	15.60	0.12	0.14
	20/09/2017	0.00	1.19	0.07	0.56
	06/10/2017	0.02	0.38	0.29	0.72
Harvested Out					

www.marine.ie

ISSN 1649-5055

www.marine.ie

HEADQUARTERS & LABORATORIES

MARINE INSTITUTE
Rinville
Oranmore
Co. Galway
Tel: +353 91 387 200
Fax: +353 91 387 201
Email: institute.mail@marine.ie

MARINE INSTITUTE REGIONAL OFFICES

MARINE INSTITUTE
Wilton Park House,
Wilton Place,
Dublin 2.
Tel: +353 1 7753900
Fax: +353 91 387201

MARINE INSTITUTE
Furnace
Newport
Co. Mayo
Tel: +353 98 42300
Fax: +353 98 42340