


Porcupine Bank *Nephrops* Grounds (FU16) 2021 UWTV Survey Report and catch scenarios for 2022

Mikel Aristegui¹, Marcin Blaszkowski¹, Jennifer Doyle¹, Sinead O'Brien¹, Imelda Hehir¹, Karl Bentley¹ and Ross Fitzgerald¹

¹ Fisheries Ecosystems Advisory Services, Marine Institute, Rinville, Oranmore, Galway, Ireland.



© Marine Institute 2021



Abstract

This report provides the results of the ninth underwater television on the 'Porcupine Bank *Nephrops* grounds' ICES assessment area; Functional Unit 16. The survey was multi-disciplinary in nature collecting UWTV and other ecosystem data. In total 71 UWTV stations were successfully completed in a randomised 6 nautical mile isometric grid covering the full spatial extent of the stock. The mean burrow density observed in 2021, adjusted for edge effect, was 0.14 burrows/m². The final krigged abundance estimate was 1018 million burrows with a CV of 5% and an estimated stock area of 7,130 km². The 2021 abundance estimate was 19% lower than in 2020. Using the 2021 estimate of abundance and updated stock data imply that catches in 2022 should be no more than 2804 tonnes, according to the EU MAP and ICES MSY approach (assuming that all catch is landed). Four species of sea-pen (*Virgularia mirabilis*, *Funiculina quadrangularis*, *Pennatula phosphorea* and the deepwater sea-pen *Kophobelemnion stelliferum*) were observed during the survey. Trawl marks were also observed on 38% of the stations surveyed.

Key words: *Nephrops norvegicus*, Porcupine Bank, stock assessment, geostatistics, underwater television (UWTV), sea-pens, benthos.

Suggested citation:

Aristegui, M., Blaszkowski, M., Doyle, J., O'Brien, S., Hehir, I., Bentley, K., Fitzgerald, R. 2021. Porcupine Bank *Nephrops* Grounds (FU16) 2021 UWTV Survey Report and catch scenarios for 2022. Marine Institute UWTV Survey report.

Introduction

Nephrops norvegicus are common around the Irish coast occurring in geographically distinct sandy/muddy areas where the sediment is suitable for them to construct their burrows. The *Nephrops* fishery in ICES sub-area 7 is extremely valuable with Irish landings in 2020 worth around €31 million at first sale. The *Nephrops* fishery on the Porcupine Bank takes place on a large area approximately 7,130 km² of complex muddy habitat between depths of between 330 – 580m. The fishery typically yields very large individual *Nephrops* that attain very high market prices relative to other fisheries around Ireland. International landings from the fishery peaked in the early 1980s around 4,000 tonnes but have shown a declining trend since then with some fluctuations (ICES, 2019). The total estimated Irish landings in 2020 were 1494 t which were likely to be worth €12.1 million.

In the recent past sustainability of the Porcupine Bank *Nephrops* stock has been a major concern. Consequently, a spatio-temporal closed area was developed and proposed by the North Western Waters Advisory Council (NWWRAC) and implemented between 1st June and 31st July in 2010-2012. Since 2013 the fishery closed for one month from 1st to 31st of May. Since 2011 a functional unit catch limit (actually landings) has also been in place as part of the TAC regulation (ICES, 2014). These measures were introduced due to negative trends in the various indicators used to assess the stock and ICES advised for a closure of the fishery in 2009 and 2010. The stock situation is known to have improved since 2010 following good recruitment. Scientific information for this area improved with the introduction of a dedicated Irish fisheries-science partnership trawl survey between 2010 and 2012 and the provision of commercial grade data by the Irish fishing industry since 2010 (ICES, 2014).

Nephrops spend a great deal of time in their burrows and their emergence behaviour is influenced by many factors; time of year, light intensity and tidal strength. Assessment methodologies, based on underwater television surveys, have been developed by ICES to provide a fishery independent estimate of stock size, exploitation status and catch advice (ICES, 2009 & 2013). Since 2012, UWTV surveys have been used to assess and provide catch scenarios and advice for this stock (ICES, 2013).

This was the ninth UWTV survey of the Porcupine Bank *Nephrops* grounds (FU16). The 2021 survey was multi-disciplinary in nature. The specific objectives are listed below:

1. To obtain 2021 quality assured estimates of *Nephrops* burrow densities from a randomised isometric grid of UWTV stations at 6 nautical mile spacing over the known spatial and bathymetric distribution of the stock (Figure 1).
2. To collect ancillary information from the UWTV footage collected at each station such as the presence of sea-pens, other macrobenthos and fish species and trawl marks on the sea bed.

This report details the final UWTV results of the 2021 survey and also documents other data collected during the survey. Operational survey details are available in the form of a survey narrative from the scientist in charge (MA). The 2021 abundance data are used to generate catch options for 2022 in line with the recommendations and procedures outlined at the 2013 ICES benchmark (ICES, 2013) and in the stock annex (ICES, 2013) and using the F_{msy} reference points proposed by FMSYREF4 (ICES, 2016).

Material and methods

A randomised isometric grid of stations at 6 nautical mile or 11.1km intervals was planned for the area. The boundary used to delineate the edge of the ground was based on VMS (vessel monitoring system) data of fishing activity between 2006-2011 targeting *Nephrops* (Lordan *et al.* 2012). The grid spacing was determined based on a time constraint of getting the survey completed within a time window of around 5-6 days. This resulted in 68 planned stations and were generated using the “spsamp1” function in the “sp” package (Pebesma & Bivand, 2005) of “R” (R Core Team, 2017). Data on bathymetry and backscatter were also available from the Irish National Seabed Survey and INFOMAR project (<http://www.infomar.ie/>). The stations ranged from 346-580m in depth with an average depth of 460m (Figure 1). Survey timing was generally standardised to June each year. In 2015 the national research vessel broke down prior to the survey and the survey was not carried out despite several attempts to get to this ground. In 2021, due to weather conditions in June, the FU16 survey was postponed and carried out between the 16th and the 23rd of August on the RV *Celtic Voyager*.

In 2021 due to technical issues with the HD camera system, all the 71 stations were surveyed using the Standard Definition (SD) camera system. SD stations were specified as: high quality, recorded to physical DVDs with a camera angle of 45°. These stations were reviewed on 15 inch (diagonal) CRT monitors – a paper based method.

The operational protocols used were those reviewed by the Workshop on the use of UWTV surveys for determining abundance in *Nephrops* stocks throughout European waters (ICES, 2007) and follow internationally agreed standards as recommended in the Manual for the *Nephrops* Underwater TV Surveys (TIMES) (Dobby *et al.*, 2021). These protocols are employed on other UWTV surveys in Irish waters and can be summarised as follows: at each station the UWTV sledge was deployed. Once stable on the seabed a 10 minute tow was recorded. Time referenced high definition image data were collected with a field of view or ‘FOV’ of 0.75 metre. Vessel position (DGPS) and position of sledge (using a USBL transponder) were recorded every 3 seconds. The navigational data were quality controlled using an “R” script developed by the Marine Institute (ICES, 2009). The USBL navigational data were used to calculate distance over ground for 93% all of stations. Corrected ship navigation data were used for 5 stations.

In line with recommendations of the Workshop on *Nephrops* Burrow Counting (WKNEPS), all scientists were trained/re-familiarised using SD image data for training material and a reference set (ICES, 2018). All counts were conducted by six trained scientists independent of each other after the survey. The numbers of *Nephrops* burrow systems were counted, where multiple burrows in close proximity which appeared to be part of a single system were counted as one. *Nephrops* activity in and out of burrows were counted and recorded for each station.

Following the recommendation of ICES Study Group on *Nephrops* Surveys (SGNEPS) the time for verified recounts was 7 minutes (ICES, 2009) but this was increased to 10 minutes for the Porcupine. This was because at the lower densities observed the relative scale of variation between minutes was higher than typical in other areas. Counting more minutes resulted in a more stable mean density estimates for each station.

Presence / absence for the occurrence of trawl marks, fish and other species were also recorded at each station. Presence / absence of sea-pen species were recorded to fulfil an OSPAR Special Request (ICES 2011).

Finally, if there was any time during each minute where counting was not possible due to sediment clouds or other reasons, this time was recorded and removed from the distance over ground calculations. The “R” quality control tool allowed for the data quality of navigation, speed, visual clarity and consistency in counts to be checked (an example is given in Figure 2).

In 2021 survey count data were screened to check for any unusual discrepancies using Lin’s Concordance Correlation Coefficient (CCC) with a threshold of 0.6. Lin’s CCC (Lin, 1989) measures the ability of counters to exactly reproduce each other’s counts on a scale of 1 to -1, where 1 is perfect concordance (i.e. a pairwise plot will have all points lying along the 1:1 line; a value of -1 would be generated by all points lying on the -1:1 line and a value of 0 indicates no correspondence at all). Lin’s CCC quality control plot of survey count data for stations 54 to 56 is shown in Figure 3. When the count data fell below the threshold of 0.6 a third review was carried out. The paired count data that passed the Lin’s CCC threshold was used in the analysis. When the paired counts did not pass the threshold an average of the three reviewers was deemed appropriate to use in the analysis.

Mean density was calculated by dividing the total number of burrow systems by the survey area observed. In 2021 the field of view of the camera at the bottom of the screen was estimated by extrapolation at 0.75m assuming that the sledge was flat on the seabed (i.e. no sinking). Occasionally the lasers were not visible at the bottom of the screen due to sinking in very soft mud, the impact of this is a minor under estimate of densities at stations where this occurred. The burrow systems in this area are relatively large and occurred at low density making the verification recounts relatively easy.

From 2012 to 2014 to account for the spatial co-variance and other spatial structuring a geo-statistical analysis of the mean and variance was carried out using SURFER Version 10.7.972 and the krigged estimation variance or coefficient of variance (CV) was carried out using the EVA: Estimation VAriance software (Petitgas and Lafont, 1997).

Since 2016, the geostatistical analysis was carried out using “RGeostats” package Version 11.1.1 (Renard D., *et al*, 2015) in “R” and is available as a separate R markdown document. The same basic steps were carried out as in previous years; construction of experimental variogram, and a model variogram (h) produced with an exponential model, a krigged grid file was created using all data points as neighbours, the same boundary used to estimate the domain area, the mean density, total burrow abundance and survey precision calculated.

For each UWTV station a temperature and depth profile was logged for the duration of each tow using a sled mounted and calibrated Seabird SBE39plus. This data will be processed at a later stage.

Results

In 2021, 71 stations were completed successfully on the Porcupine Bank (Figure 1). Figure 4 shows bubble plots of the variability between minutes for each station. At the lower densities observed the relative scale of variation between minutes was higher than typical in other areas, reflecting the patchy and low density.

A combined violin and box plot of the observed burrow densities is presented in Figure 5. This shows that median and mean burrow densities are similar in most years. The inter-quartile ranges are also similar. The mean burrow density observed in 2021, adjusted¹ for edge effect, was 0.14 burrows/m².

The final modelled density surfaces from 2012 to 2014 and 2016 to 2021 are shown as a heat map in Figure 6. The 2021 burrow surface shows a decrease in burrow densities to the south of the ground, and the usual high densities to the north. The abundance estimate derived from the krigged burrow surfaces (and adjusted for edge effect) decreased by 19% from 1264 million burrows in 2020 to 1018 million in 2021 (Figure 7 and Table 1) with an estimated area of the ground or domain area of 7,129 km². The estimation CV on the abundance was around 4.6% in 2021.

Trawl marks were observed at 26 out of the 68 surveyed stations (38%). The distribution of the various sea-pen species observed on the UWTV footage is shown in Figure 8. Three sea-pen species occur in the deep mud habitats around the coastal British Isles; *Virgularia mirabilis*, *Pennatula phosphorea* and *Funiculina quadrangularis* (Hughes, 1998). All three species were observed on footage during the 2021 survey. The distribution over time of a fourth sea-pen, *Kophobelemnon stelliferum*, is shown in Figure 9 as part of a species review in the UWTV database. This species has been recorded at the Porcupine Seabight in depths to 1600 m (Rice *et al.*, 1992).

The UWTV abundance data together with data from the fishery; landings, removals in number, and mean weight in the landings are shown (Table 2). The basis to the catch options table is given in Table 3. The catch and landings options calculated using the 2021 survey abundance estimate and updated stock data, imply that catches in 2022 should be no more than 2804 tonnes, according to ICES MSY approach (assuming that all catch is landed) (Table 4).

Discussion

This was the ninth systematic UWTV *Nephrops* survey of the Porcupine Bank. The distance from shore (~ 120 nautical miles), exposed nature of the area, the significant water depths involved (325m to 571m) and relatively large size of the area (>7100km²) presents significant logistical, technical and survey design challenges. The Marine Institutes carries out UWTV surveys in three pre-planned survey legs (of 12, 11 and 10 days). Priority was given to the Porcupine Bank which was successfully completed on the third leg in 2020, after moving the first two legs to other FUs due to weather conditions. The visibility and footage quality was

¹ Note the “adjusted” density estimates in this report are adjusted by dividing by 1.26 to take account of edge effect over estimation of area viewed during UWTV transects (see Campbell *et al.*, 2009).

normally excellent, burrow morphology and size were similar to other areas and the relatively low density meant that burrow identification was relatively easy.

The survey design, with a randomised 6 nautical mile isometric grid and fixed ground boundary, was the same as that used previously (Lordan, *et al.*, 2012). The total abundance estimate has increased. Catch scenarios for 2022 have been calculated using updated landings mean weight data from WGCSE (ICES, 2021). These data are usually estimated using the graded landings information for the fishing industry and sampling data carried out at sea by scientific observer. However, in 2020, due to low sampling levels caused by Covid-19 restrictions, sampling was adapted and mean weights estimated from catch samples. An average over the last three years (2018-2020) is used to account for the mean weight in the landings estimated in the recent past. The resulting catch advice for 2022 fishing at the F_{msy} point is a decrease by 14.8%), due to the lower abundance estimate compared to last year. Carrying out annual UWTV surveys to generate catch advice, while challenging, should be continued annually given the limited number of UWTV observations to date and evolving knowledge base on the spatial and temporal dynamics of this stock.

In addition to estimating *Nephrops* stock abundance UWTV surveys can be used to monitor the presence of certain benthic fauna (ICES, 2011). Sea-pens and burrowing megafauna communities have been included in the OSPAR list of threatened and/or declining species and habitats (OSPAR, 2010). As previously observed, all three sea-pen species which occur on mud habitat around Ireland are found on the Porcupine Bank, and also the deepwater sea-pen *Kophobelemnion stelliferum*. Data of the four sea-pen from FU16 are submitted as part of an annual ICES data call to Working Group on Deep-water Ecology (WGDEC). This dataset provides information regarding the impact of fisheries on components of the ecosystem. The occurrence of *F. quadrangularis* in particular is significant owing to its particularly vulnerability to trawl mortality. *F. quadrangularis* is largely absent from other *Nephrops* grounds around Ireland, although there are catches on groundfish surveys in areas where *Nephrops* are not commercially fished (Power and Lordan, 2012). The majority of the Porcupine Bank is fished at least once annually based on the methods described in Gerritsen, *et al.* (2013). The observation that 38% of stations showed some trawl marks is consistent with previous years.

Acknowledgments

We would like to express our sincere thanks and gratitude to Phillip Baugh (Master) and crew of the RV *Celtic Voyager*. Thanks also to the P&O Maritime IT & Instrumentation Technician Lukasz Pawlikowski who maintained the UWTV system throughout the survey. Thanks to Aodhán Fitzgerald, Rosemarie Butler Butler and Tricia McManus (RVOPs); Gordon Furey, Damian Crean, Barry Kavanagh, John Barry (P&O Maritime) for shore side support and survey logistics. Thanks to Chris Allsop (OCIS/FEAS) for supporting the UWTV database management.

Photography credits

Photograph used on the Cover have been provided by © Mikel Aristegui.

Disclaimer

Although every effort has been made to ensure the accuracy of the material contained in this publication, complete accuracy cannot be guaranteed. Neither the Marine Institute nor the author accepts any responsibility whatsoever for loss or damage occasioned, or claimed to have been occasioned, in part or in full as a consequence of any person acting or refraining from acting, as a result of a matter contained in this publication. All or part of this publication may be reproduced without further permission, provided the source is acknowledged.

References

- Campbell, N., Dobby, H., and Bailey, N. 2009. Investigating and mitigating uncertainties in the assessment of Scottish *Nephrops norvegicus* populations using simulated underwater television data. *ICES Journal of Marine Science* 66: 646–655. doi: 10.1093/icesjms/fsp046.
- Dobby, H., Doyle, J., Jónasson, J., Jonsson, P., Leocádio, A., Lordan, C., Weetman, A., and Wieland, K. 2021. ICES Survey Protocols – Manual for *Nephrops* underwater TV surveys, coordinated under ICES Working Group on *Nephrops* Surveys (WGNEPS). *ICES Techniques in Marine Environmental Sciences* Vol. 65. 44 pp. <https://doi.org/10.17895/ices.pub.8014>.
- EU. 2019. Regulation (EU) 2019/472 of the European Parliament and of the Council of 19 March 2019 establishing a multiannual plan for stocks fished in the Western Waters and adjacent waters, and for fisheries exploiting those stocks, amending Regulations (EU) 2016/1139 and (EU) 2018/973, and repealing Council Regulations (EC) No 811/2004, (EC) No 2166/2005, (EC) No 388/2006, (EC) No 509/2007 and (EC) No 1300/2008. *Official Journal of the European Union*, L 83: 117. <http://data.europa.eu/eli/reg/2019/472/oj>.
- Gerritsen, H.D., Minto, C., and Lordan, C. 2013. How much of the seabed is impacted by mobile fishing gear? Absolute estimates from Vessel Monitoring System (VMS) point data. *ICES Journal of Marine Science*. doi: 0.1093/icesjms/fst017.
- Hughes, D. J. 1998. Sea Pens and Burrowing Megafauna. An overview of dynamics and sensitivity characteristics for conservation management of marine SACs. Scottish Association for Marine Science.
- ICES 2007. Report of the Workshop on the use of UWTV surveys for determining abundance in *Nephrops* stocks throughout European waters (WKNEPHTV). ICES CM: 2007/ACFM: 14 Ref: LRC, PGCCDBS.
- ICES 2009. Report of the Study Group on *Nephrops* Surveys (SGNEPS). ICES CM 2009/LRC: 15. Ref: TGISUR.
- ICES 2011. Protocols for assessing the status of sea-pen and burrowing megafauna communities. ICES Advice 2011, Book 1, 2011.
- ICES 2013. Report of the Benchmark Workshop on *Nephrops* Stocks (WKNEPH), 25 February–1 March 2013, Lysekil, Sweden. ICES CM 2013/ACOM:45. 230 pp.

- ICES 2014. Report of the Working Group for Celtic Seas Ecoregion (WGCSE). ICES CM: 2014/ACOM:12.
- ICES 2016. EU request to ICES to provide FMSY ranges for selected stocks in ICES subareas 5 to 10. In Report of the ICES Advisory Committee, 2016. ICES Advice 2016, Book 5, Section 5.2.3.1.
- ICES. 2018. Report of the Workshop on *Nephrops* burrow counting (WKNEPS). 2-5 October 2018, Aberdeen, UK. ICES CM 2018/ EOSG:25.47 pp.
- ICES. 2021. Report of the Working Group for Celtic Seas Ecoregion (WGCSE). ICES (in draft).
- Lin, L. I. 1989. A Concordance Correlation Coefficient to Evaluate Reproducibility. *Biometrics* 45, 255-268.
- Lordan, C., Doyle, J., Dobby, H., Hehir, I. Fee, D., Allsop, C. & O'Neil, R. 2012. Porcupine Bank *Nephrops* Grounds (FU16) 2012 UWTV Survey Report and catch options for 2013. Marine Institute UWTV Survey report. <http://hdl.handle.net/10793/832>
- OSPAR, 2010. Background Document for Seapen and Burrowing megafauna communities. OSPAR Commission 2010, London. Publication number: 481/2010
- Petitgas and Lafont, 1997. EVA (Estimation VAriance). A geostatistical software on IBM-PC for structure characterization and variance computation. Version 2.
- Power, J., and Lordan, C., 2012. A review of the effects of bottom trawling on soft sediments; sea pens and burrowing megafauna biotope complexes. Marine Institute internal report Version 1, 30 July 2012.
- Renard D., Bez N., Desassis N., Beucher H., Ors F., Laporte F., 2015. RGeostats: The Geostatistical package [version:11.0.3]. MINES ParisTech. Free download from: <http://cg.ensmp.fr/rgeostats>
- Rice A.L., Tyler P.A. and Paterson G.J.L., 1992. The Pennatulid *Kophobelemnon stelliferum* (Cnidaria:Octocorallia) in the Porcupine Seabight (north-east Atlantic Ocean). *Journal of the Marine Biological Association of the UK* 72(02):417 - 434 · May 1992.
- R Core Team (2017). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>




Figure 1: FU16 UWTV 2021. UWTW map of station positions and bathymetry. The polygon line indicates the ground boundary currently used.

Source: INFOMAR is the Department of Communications, Climate Action and Environment (DCCAE) funded national seabed mapping programme, jointly managed and delivered by Geological Survey Ireland and Marine Institute




Figure 2: FU16 UWTV 2021. UWTW example quality control plot for the navigational and recount data.




Figure 3: FU16 UWTV 2021. Lin's CCC quality control plot of count data for stations 54 to 56 from the 2021 survey.




Figure 4: FU16 UWTV 2021. UWTV quality control plot showing variability between minutes for each UWTV station




Figure 5: FU16 UWTV 2021. Violin and box plot of adjusted burrow density distributions by year from 2012-2021. The blue line indicates the mean density over time. The horizontal black line represents the median, white box is the inter quartile range, the black vertical line is the range and the black dots are outliers. No UWTV survey in 2015.




Figure 6: FU16 UWTV 2021. Heat map of *Nephrops* burrow density observations from 2012 to 2021.




Figure 7: FU16 UWTV 2021. Time series of total abundance estimates for FU16 and 95% confidence intervals. No UWTV survey in 2015.




Figure 8: FU16 UWTV 2021. Stations where sea-pens *Virgularia mirabilis* (VAM), *Funiculina quadrangularis* (FAQ), *Pennatula phosphorea* (PNP) and *Kophobelemnion stelliferum* (KOP) were identified from video footage. Coloured circles denote presence in the TV station and empty circles denote TV stations with no sea-pen observations.




Figure 9: FU16 UWTV 2021. The presence/absence distribution of the deep water sea-pen species *Kophobelemnon stelliferum* (KOP) observed on the video footage from 2012 to 2021 (no survey in 2015). Coloured circles denote presence in the TV station and empty circles denote absence.

Table 1: FU16 UWTV 2021. Summary of univariate statistics and geostatistics for the burrow density estimates (bias corrected) on the Porcupine Bank UWTV survey in 2012-2021. No TV survey in 2015.

Year	Univariate Statistics						Geostatistics		
	Number of Observations	Min	Max	Mean	Standard Deviation	Variance	Adjusted abundance estimate (millions)	Domain area (km ²)	Coef. Of Variation
2012	47	0.014	0.358	0.151	0.063	0.005	787	7108	0.049
2013	68	0.012	0.233	0.106	0.051	0.003	768	7108	0.044
2014	67	0	0.226	0.099	0.049	0.002	722	7108	0.025
2015	0								
2016	65	0.011	0.325	0.132	0.0055	0.004	958	7108	0.036
2017	63	0.037	0.496	0.118	0.0082	0.006	850	7134	0.054
2018	69	0.095	0.524	0.156	0.011	0.009	1117	7130	0.042
2019	65	0.012	0.440	0.139	0.011	0.008	1010	7131	0.051
2020	65	0.0097	0.364	0.174	0.0097	0.008	1264	7133	0.038
2021	71	0	0.450	0.141	0.0096	0.008	1018	7129	0.046

Table 2: FU16 UWTV 2021. Inputs to catch scenarios table.

Year	UWTV abundance estimate	95% Confidence Interval	Landings in number	Total discards in number*	Removals in number	Harvest rate (by number)***	Landings	Total discards*	Discard proportion (by number)	Dead discard proportion (by number)	Mean weight in landings	Mean weight in discards
2012	787	79	25	0	25	3.2	1258	0	0	0	50.4	NA
2013	768	61	20	0	20	2.6	1141	0	0	0	57.5	NA
2014	722	35	17	0	17	2.4	1189	0	0	0	68.5	NA
2015	NA	NA	27	0	27	3.3**	1394	0	0	0	50.9	NA
2016	958	68	53	NA	53	5.6	2154	NA	NA	NA	40.3	NA
2017	850	90	85	NA	85	10.0	2632	NA	NA	NA	31.0	NA
2018	1117	92	66	NA	66	5.9	2751	NA	NA	NA	41.6	NA
2019	1010	101	42	NA	42	4.1	2229	NA	NA	NA	53.4	NA
2020	1263	94	49	NA	49	3.9	1877	NA	NA	NA	38.3	NA
2021	1018	92										

*Discarding up to 2015 was considered to be negligible. Discard estimates are not available since 2016 and are therefore not included in the assessment.

** The harvest rate is estimated based on a linear interpolation of abundance for 2015 as no survey was carried out in this year.

*** Values since 2016 onwards may be underestimates due to insufficient discard data.

NA = not available.

Table 3: FU16 UWTV 2021. The basis for the catch advice and scenarios.

Variable	Value	Notes
Stock abundance (2022)	1018	UWTV survey 2021; numbers of individuals in millions
Mean weight in projected landings	44.4	Average 2018–2020; in grammes
Mean weight in projected discards	-	Unknown
Projected discards	-	Discarding assumed negligible
Discards survival	-	Not applicable
Projected dead discards	-	Assumed to be zero

Table 4: FU16 UWTV 2021. Annual catch advice and scenarios.

Basis	Total catch	Projected landings	Projected discards	Harvest rate * %	% advice change **
	PL + PD	PL	PD	For PL + PD	
ICES advice basis					
EU MAP [^] : F _{MSY}	2804	2804	0	6.2	-14.8
EU MAP [^] : F _{MSY lower}	2261	2261	0	5.0	-14.8
EU MAP [^] : F _{MSY upper} ***	2804	2804	0	6.2	-14.8
Other scenarios					
MSY approach	2804	2804	0	6.2	-14.8
F ₂₀₂₀	1755	1755	0	3.9	-47

* By number

** Advice values for 2022 relative to the corresponding 2021 values (MAP advice of 3290, 2653, and 3290 tonnes, respectively); other option values are relative to 3290 t.

*** F_{MSY upper} = F_{MSY} for this stock

[^] EU multiannual plan (MAP) for Western Waters (EU, 2019)