

The Labadie, Jones and Cockburn Banks *Nephrops* Grounds (FU2021) 2020 UWTV Survey Report and catch scenarios for 2021.

Mikel Aristegui¹, Dave Tully¹, Marcin Blaszkowski¹, Jennifer Doyle¹, Dermot Fee¹, Seán O'Connor¹ and Jonathan White¹

¹ Fisheries Ecosystems Advisory Services, Marine Institute, Renville, Oranmore, Galway, Ireland.



Version 1.0, October 2020

Abstract

This report provides the main results of the 2020 underwater television survey on the 'Labadie, Jones and Cockburn Banks' ICES assessment area; Functional Unit 2021. The 2020 survey was multi-disciplinary in nature collecting UWTV, and other ecosystem data. A total of 97 UWTV stations were completed at 6nm intervals over a randomised isometric grid design. The mean burrow density was 0.102 burrows/m² compared with 0.06 burrows/m² in 2019. The 2020 geostatistical abundance estimate was 1020 million, a 65% increase on the abundance from 2019, with a CV of 5%, which is well below the upper limit of 20% recommended by SGNEPS 2012. Low to medium densities were observed throughout the ground. Using the 2020 estimate of abundance and updated stock data implies catch in 2021 that correspond to the F ranges in the EU multi annual plan for Western Waters are between 1682 and 1710 tonnes (assuming that discard rates and fishery selection patterns do not change from the average of 2017–2019). One species of sea-pen (*Virgularia mirabilis*) were recorded as present at the stations surveyed. Trawl marks were observed at 36% of the stations surveyed.

Key words: *Nephrops norvegicus*, Celtic Sea, stock assessment, geostatistics, underwater television (UWTV), benthos.

Suggested citation:

Aristegui, M., Blaszkowski, M., Doyle, J., Fee, D., O'Connor, S. and White, J., 2020. The Labadie, Jones and Cockburn Banks *Nephrops* Grounds (FU2021) 2020 UWTV Survey Report and catch scenarios for 2021. Marine Institute UWTV Survey report.

Introduction

The prawn (*Nephrops norvegicus*) are common in the Celtic Sea occurring in geographically distinct sandy/muddy areas where the sediment is suitable for them to construct their burrows. The *Nephrops* fishery in ICES sub-area 7 is extremely valuable with Irish landings in 2019 worth around €42 million at first sale. The Celtic Sea area (Functional Units 19-22; Figure 1) support a large multi-national targeted *Nephrops* fishery, mainly using otter trawls and yielding landings in the region of ~5,000 t annually over the last decade (ICES, 2018). The 2019 reported landings from FU2021 were ~2,219 t and estimated to be worth in the region of €12 million at first sale. This ground has become increasingly important to the Irish demersal fleet, which now accounts for around 74% of the total international FU2021 *Nephrops* landings (ICES, 2019). Good scientific information on stock status and exploitation rates are required to inform sustainable management of this resource.

Nephrops spend a great deal of time in their burrows and their emergence behaviour is influenced by several factors: time of day, time of year, light intensity, tidal strength, etc. Underwater television surveys and assessment methodologies have been developed to provide a fishery independent estimate of stock size, exploitation status and catch advice (ICES, 2009a & 2012a). This is the eighth UWTV survey in the Celtic Sea FU2021 grounds carried out by the Marine Institute, Ireland.

The survey was multi-disciplinary in nature and the 2020 specific objectives are listed below:

1. To complete 98 UWTV stations with 6.0 nautical mile (nm) spacing on the Labadie, Jones and Cockburn *Nephrops* ground (FU2021).
2. To obtain 2020 quality assured estimates of *Nephrops* burrow distribution and abundance, and compare them with those from previous surveys.
3. To collect ancillary information from the UWTV footage collected at each station such as the occurrence of sea-pens, other macro benthos and fish species and trawl marks on the sea bed.
4. To collect sediment samples for PSA analysis.

This report details: the survey design, the final UWTV results of the 2020 survey and also documents other data collected during the survey. Operational survey details are available in the form of a survey narrative from the scientist in charge (MA). The 2020 abundances are used to generate catch advice scenarios for 2021 in line with the recommendations and procedures outlined in the stock annex for FU2021 (ICES, 2020).

Material and methods

The knowledge about the distribution of suitable *Nephrops* habitat in this area has been developing. Information so far suggests that *Nephrops* are found in complex channels, which are probably the remnants of fluvial channels related to the deglaciation of the Irish ice sheet at the end of the last ice age. The area of the ground was revised by WKCELT (ICES, 2014) with the inclusion of both French and Irish integrated logbook VMS data (Gerritsen & Lordan, 2011) and is now calculated at 10,014 km². This value is used for the survey.

INFOMAR seabed mapping programmes are now focussed in the Celtic Sea and the multibeam backscatter data from these surveys will aid in any area redefinition in time.

INFOMAR is the Department of Communications, Climate Action and Environment (DCCA) funded national seabed mapping programme, jointly managed and delivered by Geological Survey Ireland and Marine Institute. Figure 2 shows the backscatter data to date relevant to this *Nephrops* survey area, where light grey indicates soft sediment and darker grey harder ground.

The 2020 randomised isometric grid resulted in 98 planned stations and was generated using the “spsamp1” function in the “sp” package (Pebesma & Bivand, 2005) in “R” (R Core Team, 2017). Station depths varied from 73m to 140m and the completed stations ranged from 30 to 134 nm offshore (Figure 3). In 2020, FU2021 was surveyed during two surveys onboard the RV. Celtic Voyager: the first one from the 20th of July to the 5th of August and the second one from the 8th to the 16th of August.

In 2020 image data were collected by a custom built camera system recording High Definition still image data at 12 frames per second with a camera angle of 75°. The digital images were stored on a server and were reviewed after the survey through an inhouse developed Image annotation R Shiny app (Aristegui, 2020). This app allows each reviewer to annotate burrows for each randomly assigned station in an efficient manner. The survey process onboard is now paperless.

The operational protocols used were those reviewed by WKNEPHTV 2007 (ICES, 2007) and employed on other UWTV surveys in Irish waters. These protocols can be summarised as follows: At each station the UWTV sledge was deployed. Once stable on the seabed a 10 minute tow was recorded. Time referenced high definition image data with field of view or ‘FOV’ of 1.01 metre. Vessel position (DGPS) and position of sledge (using a USBL transponder) were recorded every 3 seconds. The navigational data was quality controlled using an “R” script developed by the Marine Institute (ICES, 2009b). In 2020 USBL navigational data were used to calculate distance over ground or ‘DOG’ for all the stations.

In line with recommendations of the Workshop on *Nephrops* Burrow Counting (WKNEPS), all scientists were trained/re-familiarised using 2019 image data for training material and reference set (ICES, 2018a). All counts were conducted by six trained scientists independent of each other after the survey.

During this review numbers of *Nephrops* burrows systems were counted, where multiple burrows in close proximity which appear to be part of a single system were counted as one. *Nephrops* activity in and out of burrows were counted and recorded for each station. Following the recommendation of SGNEPS the time for verified recounts was 7 minutes (ICES, 2009a).

Presence / absence notes were also recorded each station on the occurrence of trawl marks, fish and other species. Presence / absence of sea-pen species were also recorded to fulfil an OSPAR Special Request (ICES 2011). Finally, if there was any time during the one-minute where counting was not possible, due to sediment clouds or other reasons, this was also estimated so that the time window could be removed from the distance over ground calculations. The “R” quality control tool allowed the quality of data for each station to be checked: navigation, speed, visual clarity and consistency in counts (Figure 4).

In 2020 the survey count data were screened to check for any unusual discrepancies using Lin's Concordance Correlation Coefficient (CCC) with a threshold of 0.5. Lin's CCC (Lin, 1989) measures the ability of counters to exactly reproduce each other's counts on a scale of 1 to -1 where 1 is perfect concordance (i.e. a pairwise plot will have all points lying along the 1:1 line, a value of -1 would be generated by all points lying on the -1:1 line and a value of 0 indicates no correspondence at all). For those stations that did not pass the threshold it was deemed appropriate to carry out a third review. The paired count data that passed the threshold were used in the analysis. When the paired counts did not pass it was deemed acceptable to use the average of the three reviewers in the analysis. Lin's CCC quality control plots of count data for stations 201, 202 and 204 are shown in Figure 5. There is moderate variability between counters but no obvious bias or excessive deviations. The moderate variability between counters is because burrow counting in this area is particularly difficult (see discussion).

Mean density was calculated by dividing the total number of burrow systems by the survey area observed. The USBL data were used to calculate distance over ground of the sledge. The field of view of the camera at the bottom of the screen was estimated by extrapolation at 1.01 metre assuming that the sledge was flat on the seabed (i.e. no sinking). Occasionally the lasers were not visible at the bottom of the screen due to sinking in very soft mud, the impact of this is a minor under estimate of densities at stations where this occurred.

For each UWTV station a temperature and depth profile was logged for the duration of each tow using a sled mounted and calibrated Seabird SBE39plus. This data will be processed at a later stage inhouse and is considered an emerging time series.

17 sediment samples were collected using the Shipex grab after completion of a TV station and when survey time allowed. Three of these samples will be used in a microplastics project by the Galway-Mayo Institute of Technology (GMIT). The other 14 samples will be used to generate sediment maps for this area and also to ground through INFOMAR sea-bed mapping programmes in this area.

The approach to work up the abundance estimates each year has been documented in previous survey reports. Since 2015 the geostatistical analysis was carried out using the "RGeostats" package (Renard D., *et al*, 2015) and is available as an "R" markdown document. The same steps were carried out as in previous years; construction of experimental variogram, a model variogram produced with an exponential model, create krigged grid file using all data points as neighbours, same boundary used to estimate the domain area, mean density, total burrow abundance and calculate survey precision.

Results

97 out of the 98 planned stations were completed successfully on the FU2021 *Nephrops* grounds (Figure 3), and only one station was dropped due to fishing gear on the area. Figure 6 shows bubble plots of the variability between minutes and operators. These show that the burrow estimates are fairly consistent between minutes and counters. The variability is slightly higher between minutes.

The adjusted burrow densities from 2013 to 2020 are shown in Figure 7 as a combined violin and box plot. These show that density has increased in 2020 from 2019. The highest station density observed in 2020 was 0.34 burrows/m² while the majority were in the lower range of < 0.15 burrows/m². The 2020 mean adjusted¹ density of 0.10 burrows/m² is the second lowest in the time series to date, but was 67% higher than the 2019 estimate of 0.06 burrows/m².

Combined bubble and contour plots of the krigged densities from 2013 to 2020 are presented in Figure 8. Highest densities were towards the centre of the ground in years 2013 - 2014 while for 2015 - 2016 high densities were found in the northern and south-western area. In 2017 high densities were generally observed throughout the ground but the highest were to the northwest. In 2018 high densities were observed throughout the ground. There were also high densities observed close to boundaries in several areas. There was a general decrease in densities observed in 2019, which slightly increased in 2020.

The summary empirical and geo-statistical results are given in Table 1. Stations surveyed in 2006 and 2012 should be viewed as exploratory and have not been used to extrapolate total abundance. The 2013 survey had partial coverage of the area (<60%) scaling the mean density to the total area (10,014 km²) resulted in an abundance estimate of 1.6±0.3 billion. The 2020 geo-statistical abundance estimate is 1020 million, which is 65% higher than in 2019 (Figure 9). The geo-statistical CVs were always in the order of 3 to 5%, which are well below the upper limit recommended of 20% (ICES, 2012).

The UWTV abundance data together with data from the fishery; landings, discards and removals in number are used to calculate the harvest rate (dead removals/TV abundance) in 2019, which was 21.2% (Table 2). The mean weight in the landings and the discards and the proportions of removal retained are also shown (Table 2). The mean weights are variable between 2012 – 2019 and are based on available sampling data (ICES, 2019). The basis to the catch scenarios is given in Table 3. The catch scenarios and associated harvest rates are presented in Table 4. When the EU multiannual plan (MAP) for Western Waters and adjacent waters is applied (EU, 2019), catches in 2020 that correspond to the F ranges in the MAP are between 1682 and 1710 t (Table 4). This assumes that discard rates and fishery selection patterns do not change from the average of 2017–2019.

Sea-pen distribution across the *Nephrops* grounds is mapped in Figure 10. All sea-pens were identified from the image data as *Virgularia mirabilis*. Trawl marks were noted at 36% of the stations surveyed.

Discussion

The 2020 survey achieved full coverage of the stock area for the seventh time. The density estimates in 2013 – 2016 are relatively similar and would be considered low (mainly ~0.2m²). In 2017 there was a large increase in the densities across the whole ground, with a large increase also observed in FU22 also in the Celtic Sea (O'Brien *et al.*, 2017). In 2018 and 2019

¹ Note the “adjusted” density estimates in this report are adjusted by dividing by 1.3 (Table 2) to take account of edge effect over estimation of area viewed during UWTV transects (see Campbell *et al* 2009).

density across FU2021 decreased. There was also a general decrease in density in the Celtic Sea *Nephrops* grounds in FU19 (Aristegui *et al.*, 2018a) and FU22 (Aristegui *et al.*, 2018b). In 2020 density across FU2021 increased slightly, but it still the second lowest abundance estimate of the time series. Fluctuations in density were also observed in the adjacent FU19 and FU22 in recent years (Doyle *et al.*, 2019a; Doyle *et al.*, 2019b; Aristegui *et al.*, 2020b; Aristegui *et al.*, 2020a). Sudden declines followed by large increases in abundance have also been observed in other *Nephrops* stocks in the past (e.g. FU12 and FU13 in 2012-2013).

In 2018, a partial review of historical survey data was undertaken given the large fluctuations observed in the short time series to date for this survey, that is, to randomly check 20% of UWTV stations in years 2016 and 2017 as recommended by the working group for the Celtic Seas Ecoregion WGCSE (ICES, 2018b). This process was conducted in July 2018 during the FU2021 UWTV survey. The analysis was presented to WGNPS where full details are available in R-markdown (ICES, 2018c; Annex 7). This process was also followed in 2019 given the steep decrease in abundance estimate and following guidelines set out in the Manual for the *Nephrops* Underwater TV Surveys (SISP) (ICES, in prep). The analysis showed that the 2019 observed densities observed are robust (ICES, 2019).

It is likely that the recruitment to *Nephrops* populations in the Celtic Sea are linked through oceanographic process (O'Sullivan *et. al.*, 2015). It may well be that oceanographic conditions have resulted in the observed density decreases in the Celtic Sea.

Nephrops fisheries in this area have been covered under the landings obligation since 2016 with several exemptions. Discard rates in weight for this FU have been around 16% in recent years.

The provision of catch advice and scenarios for 2021 based on the EU multiannual plan (MAP) for Western Waters assumes that discard rates and fishery selection patterns do not change from the average of 2017–2019.

The introduction of the landings obligation to *Nephrops* fisheries in 2016 should result in changes in selectivity. This is not taken into account in any of the catch advice because it is not possible to predict exactly what might happen. The main message is that any improvements in selectivity in the fishery and reductions in discards will result in increased mean weight in the catches. This will in turn reduce overall mortality on the stocks and allow for catch increases in the future.

An important objective of this UWTV survey is to collect various ancillary information. The occurrence of trawl marks on the footage is notable for two reasons. Firstly, it makes identification of *Nephrops* burrows more difficult as the trawl marks remove some signature features making accurate burrow identification more difficult. Secondly, only occupied *Nephrops* burrows will persist in heavily trawled grounds and it is assumed that each burrow is occupied by one individual *Nephrops* (ICES 2008).

Monitoring the occurrence and frequency of sea-pens observed on these *Nephrops* patches is important in the context of OSPAR's designations of sea-pen and burrowing megafauna communities as threatened. The sea-pen species *Virgularia mirabilis* which was seen in 2020

have been observed on previous surveys of FU2021. Monitoring *Nephrops* stock and the benthic habitat is also important in the context of the MFSD indicators (e.g. sea floor integrity).

The sediment sample data collected during the survey will increase the knowledge base on habitat mapping in time.

The main objectives of the survey were successfully met for the seventh successive year. The UWTV coverage and footage quality was excellent throughout the survey. The multi-disciplinary nature of the survey means that the information collected is highly relevant for a number of research and advisory applications.

Acknowledgments

We would like to express our thanks and gratitude to Phillip Baugh (Master) and crew of the RV. Celtic Voyager for their good will and professionalism throughout the survey. Thanks also to the P&O Maritime IT & Instrumentation Technician Damian Crean who maintained the UWTV system throughout the survey. Thanks to Aodhán Fitzgerald, Rosemarie Butler and Tricia McManus (RVOPs) and Dave Tully FEAS at the Marine Institute for organising survey logistics. Thanks to Gordon Furey, Damian Crean, Barry Kavanagh, John Barry (P&O Maritime) for shore side support.

References

- Aristegui, M. (2020) Image annotation R Shiny app. Marine Institute. <http://doi.org/d24n>
- Aristegui, M., Blaszkowski, M., Doyle, J., Ryan, G., and McAuliffe, M. 2020a. The “Smalls” *Nephrops* Grounds (FU22) 2020 UWTV Survey Report and catch scenarios for 2021. Marine Institute UWTV Survey report.
- Aristegui, M., Doyle, J., O’Brien, S., Tully, D., McAuliffe, M., Fitzgerald, R., Fee, D., O’Connor, S., Galligan, S., Blaszkowski, M., Butler, R., and White., J. 2020b. FU19 *Nephrops* Grounds (FU19) 2020 UWTV Survey Report and catch scenarios for 2021. Marine Institute UWTV Survey report.
- Aristegui, M., O’Brien, S., Blaszkowski, M., O’Connor, S., Fitzgerald, R., and Doyle, J. 2018. FU19 *Nephrops* Grounds (FU19) 2018a UWTV Survey Report and catch scenarios for 2019. Marine Institute UWTV Survey report.
- Aristegui, M., O’Brien, S., Blaszkowski, M., O’Connor, S., Fitzgerald, R., and Doyle, J. 2018. The “Smalls” *Nephrops* Grounds (FU22) 2018b UWTV Survey Report and catch scenarios for 2019. Marine Institute UWTV Survey report.
- Campbell, N., Dobby, H., and Bailey, N. 2009. Investigating and mitigating uncertainties in the assessment of Scottish *Nephrops norvegicus* populations using simulated underwater television data. ICES Journal of Marine Science 66: 646–655. doi: 10.1093/icesjms/fsp046.

- Doyle, J., Aristegui, M., O'Brien, S., Lynch, D., Vacherot, JP., and Fitzgerald, R. 2019. FU19 *Nephrops* Grounds (FU19) 2019a UWTV Survey Report and catch scenarios for 2020. Marine Institute UWTV Survey report.
- Doyle, J., O'Brien, S., Fitzgerald, R., Vacherot, JP., Sugrue, S., and Quinn M. 2019. The "Smalls" *Nephrops* Grounds (FU22) 2019b UWTV Survey Report and catch scenarios for 2020. Marine Institute UWTV Survey report.
- EU 2019. Regulation (EU) 2019/472 of the European Parliament and of the Council of 19 March 2019 establishing a multiannual plan for stocks fished in the Western Waters and adjacent waters, and for fisheries exploiting those stocks, amending Regulations (EU) 2016/1139 and (EU) 2018/973, and repealing Council Regulations (EC) No 811/2004, (EC) No 2166/2005, (EC) No 388/2006, (EC) No 509/2007 and (EC) No 1300/2008. Official Journal of the European Union, L 83: 1–17. <http://data.europa.eu/eli/reg/2019/472/oj>
- Gerritsen, H., and Lordan, C. 2011. Integrating vessel monitoring systems (VMS) data with daily catch data from logbooks to explore the spatial distribution of catch and effort at high resolution. ICES Journal of Marine Science: Journal du Conseil 68: 245–252. doi: <http://dx.doi.org/10.1093/icesjms/fsq137>.
- ICES 2007. Report of the Workshop on the use of UWTV surveys for determining abundance in *Nephrops* stocks throughout European waters (WKNEPHTV). ICES CM: 2007/ACFM: 14 Ref: LRC, PGCCDBS.
- ICES 2008. Report of the Workshop and training course on *Nephrops* burrow identification (WKNEPHBID). ICES CM: 2008/LRC: 3 Ref: LRC, ACOM.
- ICES 2009a. Report of the Study Group on *Nephrops* Surveys (SGNEPS). ICES CM 2009/LRC: 15. Ref: TGISUR.
- ICES 2009b. Report of the Benchmark Workshop on *Nephrops* assessment (WKNEPH). ICES CM: 2009/ACOM:33
- ICES 2011. Report of the ICES Advisory Committee 2011. ICES Advice.2011. Book 1: Introduction, Overviews and Special Requests. Protocols for assessing the status of sea-pen and burrowing megafauna communities, section 1.5.5.3.
- ICES 2012. Report of the Study Group on *Nephrops* Surveys (SGNEPS). ICES CM 2012/SSGESST: 19. Ref: SCICOM, ACOM
- ICES 2014. Report of the Benchmark Workshop on Celtic Sea stocks (WKCELT), 3–7 February 2014, ICES Headquarters, Copenhagen, Denmark. ICES CM 2014\ACOM:42. 194 pp.
- ICES 2016b. EU request to ICES to provide FMSY ranges for selected stocks in ICES subareas 5 to 10. In Report of the ICES Advisory Committee, 2016. ICES Advice 2016, Book 5, Section 5.2.3.1.
- ICES 2018a. Report of the Workshop on *Nephrops* burrow counting, 2-5 October 2018, Aberdeen, UK. ICES CM 2018/ EOSG:25.47 pp.
- ICES 2018b. Report of the Working Group for Celtic Seas Ecoregion (WGCSE). ICES CM: 2018/ACOM:13 in draft.

- ICES 2018c. Report of the Working Group on *Nephrops* Surveys (WGNEPS), 6–8 November 2018. Lorient, France. ICES CM 2018/ EOSG:18.227 pp.
- ICES 2020. Report of the Working Group for Celtic Seas Ecoregion (WGCSE). ICES (in draft).
- ICES (in prep.). Manual for the *Nephrops* Underwater TV Surveys. Series of ICES Survey Protocols (SISP).
- Lin, L. I. 1989. A Concordance Correlation Coefficient to Evaluate Reproducibility. *Biometrics* 45, 255-268.
- O’Sullivan D, Lordan C, Doyle J, Berry A, Lyons K. 2015. Metapopulation connectivity via larval transport of the Norway lobster *Nephrops norvegicus* in waters around Ireland: a modelled approach. *Mar Ecol Prog Ser* 534:95-106.
- O’Brien, S., Blaszkowski, M., Butler, R., Fee, D., Hernon, P., Santana, C., Lordan, C. and Doyle, J. 2017. The “Smalls” *Nephrops* Grounds (FU22) 2017 UWTV Survey Report and catch options for 2018. Marine Institute UWTV Survey report.
- Pebesma, E.J., R.S. Bivand, 2005. Classes and methods for spatial data in R. *R News* 5 (2), <https://cran.r-project.org/doc/Rnews/>
- R Core Team 2017. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>
- Renard D., Bez N., Desassis N., Beucher H., Ors F., Laporte F. 2015. RGeostats: The Geostatistical package [version:11.1.1]. MINES ParisTech. Free download from: <http://cg.ensmp.fr/rgeostats>




Figure 1: FU2021 UWTV 2020. *Nephrops* Functional Units (FUs) and *Nephrops* survey area polygons in the greater Celtic Sea.




Figure 2: FU2021 UWTV 2020. Multibeam backscatter data from INFOMAR mapping programme to date. Green (+) are UWTV stations completed to date and green shading is the extent of UWTV survey area.




Figure 3: FU2021 UWTW 2020. UWTW stations on the 2020 survey.




Figure 4: FU2021 UWTW 2020. R - tool quality control plot for station 193 of the 2020 survey.




Figure 5: FU2021 UWTV 2020. Lin's CCC quality control plots of count data for stations 201, 202, 204 from the 2020 survey.




Figure 6: FU2021 UWTV 2020. Plots of the variability in density between minutes for each station in 2020.




Figure 7: FU2021 UWTV 2020. Violin and box plot of adjusted burrow density distributions by year from 2013-2020. The blue line indicates the mean density over time. The horizontal black line represents medians, white boxes the inter quartile ranges, the black vertical lines are the range and the black dots are outliers.




Figure 8: FU2021 UWTV 2020. Contour plots of the krigged density estimates by year from 2013 (top-left) - 2020 (bottom-right).




Figure 9: FU2021 UWTV 2020. Time series of raised abundance estimates (in millions of burrows) for FU2021. The error bars indicate the 95% confidence intervals.




Figure 10: FU2021 UWTV 2020: 2020 stations where *Virgularia mirabilis* (VAM) were identified. Pink circles indicated presence and open circles denotes TV stations with no sea-pen observations.

Table 1: Summary of UWTV results; number of stations, mean density observed, standard deviation, absolute abundance estimates with 95% confidence intervals, estimated area of the stock and coefficient of variation on the abundance.

Year	Number of stations	Mean Density adjusted (burrow /m ²)	Standard Deviation	Absolute abundance estimate (million burrows)	95 % CI on Abundance	Domain area	CVs %
2006	9	0.44	0.31	nr			
2012	54	0.57	0.25	nr			
2013	55	0.16	0.11	942	60	5701	3
2013*				1624	103	9835	
2014	98	0.19	0.14	2051	131	9835	3
2015	96	0.20	0.02	2003	118	9835	3
2016	93	0.18	0.02	1879	147	9835	5
2017	86	0.44	0.08	4428	347	9835	4
2018	96	0.27	0.04	2721	212	9835	4
2019	95	0.06	0.004	617	58	9835	5
2020	97	0.10	0.009	1020	96	9835	5

* the 2013 survey achieved partial coverage ~60% of the total area. The abundance has been scaled up to the entire area since densities in the un-surveyed part of the ground were not significantly different in 2014.

nr = no reliable abundance estimate could be calculated because survey coverage was partial.

Table 2: The inputs for the catch scenarios.

Year	UWTV abundance estimate	95% Confidence Interval	Landings in number	Total discards in number*	Removals in number	Harvest rate (by number)	Landings	Total discards*	Discard proportion (by number)	Dead discard proportion (by number)	Mean weight in landings	Mean weight in discards
	Millions				%	tonnes	%		grammes			
2012			38.2	36.1	65.3		1189	542	48.5	41.4	31.1	15.0
2013	1624	103	34.8	19.2	49.2	3.0	1387	327	35.6	29.3	39.9	17.0
2014	2051	131	50.6	55.5	92.2	4.5	1836	834	52.3	45.2	36.3	15.0
2015	2003	129	59.4	28.1	80.5	4.0	2116	442	32.2	26.2	35.7	15.7
2016	1879	157	60.2	37.5	88.3	4.7	2453	801	38.4	31.8	40.7	21.4
2017	4428	145	60.1	19.2	74.5	1.7	1849	306	24.3	19.4	30.8	15.9
2018	2721	212	64.7	21.5	80.8	3.0	1803	381	25.0	20.0	27.9	17.7
2019	617	58	99.1	42.3	130.9	21.2	2999	637	29.9	24.3	30.2	15.0
2020	1020	96										

Table 3: The basis for the catch scenarios.

Variable	Value	Notes
Stock abundance (2021)	1020	Numbers of individuals (millions); UWTV survey 2020
Mean weight in projected landings	29.6	Average 2007–2019 in grammes
Mean weight in projected discards	16.2	Average 2007–2019 in grammes
Projected discards	26.4	Proportion by number; Average 2017–2019
Discards survival	25	Proportion by number
Projected dead discards	21.2	Proportion by number; Average 2017–2019

Table 4: Catch advice and scenarios for 2020; Discarding assumed to continue at recent average. All weights are in tonnes.

Basis	Total catch	Dead removals	Projected landings	Projected dead discards	Projected surviving discards	Harvest rate* %	% advice change **
	PL + PDD + PSD	PL + PDD	PL	PDD	PSD	for PL + PDD	
ICES advice basis							
EU MAP [^] : F _{MSY}	1710	1640	1430	211	70	6.0	48.7
F = MAP F _{MSY lower}	1682	1613	1406	207	69	5.9	48.7
F = MAP F _{MSY upper} ***	1710	1640	1430	211	70	6.0	48.7
Other options							
MSY approach	1710	1640	1430	211	70	6.0	48.7
F ₂₀₁₈	6048	5800	5056	745	248	21.2	425.9

[^] EU multiannual plan (MAP) for Western Waters (EU, 2019).

* By number.

** Advice value for 2021 relative to the corresponding 2020 values (MAP advice value of 1150, 1131 and 1150 tonnes respectively; other values are relative to F_{MSY}).

*** F_{MSY upper} = F_{MSY} for this stock